



stats sa

Department:
Statistics South Africa
REPUBLIC OF SOUTH AFRICA

Private Bag X44, Pretoria, 0001, South Africa, ISibalo House, Koch Street, Salvokop, Pretoria, 0002

www.statssa.gov.za, info@statssa.gov.za, Tel +27 12 310 8911

STATISTICAL RELEASE

P0309.3

Mortality and causes of death in South Africa: Findings from death notification 2024

Embargoed until:

22 September 2026

11:30

ENQUIRIES:
User information services
012 310 8600

FORTHCOMING ISSUE:
2025

EXPECTED RELEASE DATE
TO BE CONFIRMED

Dipalopalo tsa Aforikaborwa • Dipalopalo tsa Aforika Borwa • Ezazibalo zaseNingizimu Afrika • Tshitatistika Afrika Tshipembe • Tinhlayo Afrika-Dzonga

Statistieke Suid-Afrika • Dipalopalo tša Aforika Borwa • Telubalo zaseNingizimu Afrika • EzeeNkcukacha maNani zoMzantsi Afrika • Iimbalobalo zeSewula Afrika

IMPROVING LIVES THROUGH DATA ECOSYSTEMS



Preface

This statistical release presents information on mortality and causes of death in South Africa for deaths that occurred during the 2024 calendar year. Deaths that occurred during the years 2003–2023 have also been included to show trends in mortality and causes of death, using updated information which includes late registrations. The statistical release is based on deaths collected through the South African civil registration system maintained by the Department of Home Affairs. The information on causes of death is captured as it was recorded on death notification forms completed by medical practitioners and other certifying officials.



Risenga Maluleke

Statistician-General

Table of contents

Preface	i
Table of contents	ii
List of tables	iv
Abbreviations/acronyms	vi
1. Introduction	1
1.1 Background	1
1.2 Objectives of this statistical release	2
1.3 Scope of this statistical release	2
1.4 Organisation and presentation of this statistical release	2
2. Data and methods	3
2.1 Data source	3
2.2 Data processing	4
2.2.1 Classification of the causes of death	4
2.2.2 Generation of the underlying causes of death	5
2.3 Data editing	5
2.4 Assessment of the quality of data	6
2.5 Data analysis	6
3. Mortality	8
3.1 Levels and trends of mortality in 2023	8
3.2 Age differentials	10
3.3 Sex differentials	12
3.4 Age and sex differentials	14
3.4.1 Distribution of deaths by age group and sex	14
3.4.2 Median ages at death by sex	16
3.4.3 Sex ratios by age groups	18
3.5 Population group differences in mortality	20
3.6 Marital status differences in mortality	20
3.7 Differences in mortality by smoking status of the deceased	21
3.8 Differences in mortality by place or institution of death occurrence	22
3.9 Geographic variations in mortality	22
3.9.1 Differences by province, age and sex	22
4. Causes of death	24
4.1 Reported causes of death	24
4.2 Method of ascertaining cause of death	25
4.3 Main groups of the underlying causes of death	27
4.4 Natural and non-natural causes of death	29
4.4.1 Natural and non-natural causes of death by age	31
4.5 Major groups of causes of death as per Global Burden of Disease	33
4.6 Broad groups of natural causes of death	38

4.6.1 Overall pattern of the leading underlying natural causes of death	38
4.6.2 Leading underlying natural causes of death by sex	39
4.6.3 Leading underlying natural causes of death by age	42
4.6.4 Leading underlying natural causes of death for children aged below five years by age groups ...	44
4.6.5 Leading underlying natural causes of death for the population aged 15–24 years	46
4.6.6 Leading underlying natural causes of death by province of death occurrence	46
4.7 Non-natural causes of death	49
4.7.1 Non-natural causes of death by age and sex	51
4.7.2 Non-natural causes of death by province of death occurrence	54
4.8 Comparison between immediate, contributing and underlying causes of death	56
5. Conclusion	59
6. References	61
Appendices	63
Appendix A: Glossary	63
Appendix B: Death Notification form	64
Appendix C: Assessment of the quality of data	64

List of tables

Table 3.1 - Number and percentage (%) distribution of deaths by age group, 2023	10
Table 3.2 - Number and percentage (%) distribution of deaths by population group, 2023.....	20
Table 3.3 - Number and percentage (%) distribution of deaths by marital status, 2023	21
Table 3.4 - Number and percentage (%) distribution of deaths by smoking status among those aged 16 and older, 2023	21
Table 3.5 - Number and percentage (%) distribution of deaths by place of death occurrence, 2023	22
Table 3.6 - Distribution of deaths by province of death occurrence and province of usual residence of the deceased, 2023	23
Table 4.1 - Number and percentage (%) distribution of death notification forms by the number of causes reported on the death notification form, 2023	25
Table 4.2 - Number and percentage (%) distribution of deaths by method used to ascertain the cause of death, 2023	26
Table 4.3 - Distribution of deaths by main causes of death, 2023	28
Table 4.4 - Number of natural and non-natural deaths by year of death occurrence, 2002–2023*	30
Table 4.5 – The ten leading underlying natural causes of death, 2021–2023*	39
Table 4.6 - The ten leading underlying causes of death for males and females, 2023*	40
Table 4.7 - The ten leading underlying natural causes of death for broad age groups, 2023.....	43
Table 4.8 - The ten underlying natural causes of death for infants and children aged below five years, 2023 ..	45
Table 4.9 - The ten leading underlying causes of death for the population aged 15–24 years, 2023	46
Table 4.10 - The ten leading underlying natural causes of death in each province of death occurrence, 2023 ..	48
Table 4.11 - Distribution of non-natural causes of death by broad groups, 2023	49
Table 4.12 - Distribution of deaths due to other external causes of accidental injury, 2023.....	50
Table 4.13 - Underlying non-natural causes of death by age group and sex, 2023	53
Table 4.14 - Underlying non-natural causes of death by province, 2023.....	55
Table 4.15 - Distribution of the 20 most commonly reported causes of death, 2023.....	57
Table 4.16 - Number and percentage (%) of deaths selected as underlying or reported as immediate or contributing causes of death, 2023.....	58

List of figures

Figure 3.1 - Number of registered deaths by year of death, 2002–2023*	9
Figure 3.2 - Percentage (%) distribution of deaths by age group and year of death, 2019–2023*	11
Figure 3.3 - Percentage (%) distributions of deaths by sex and year of death, 2002–2023*	12
Figure 3.4 - Percentage (%) distribution of deaths by age group and sex, 2023*	15
Figure 3.5 - Median ages at death by sex and year of death, 2002–2023*	17
Figure 3.6 - Sex ratios by age group and year of death, 2019–2023*	19
Figure 4.1 - Percentage (%) distribution of deaths by main groups of causes of death, 2021–2023*	29
Figure 4.2 - Percentage (%) distribution of natural and non-natural causes of death by year of death, 2002–2023*	31
Figure 4.3 - Percentage (%) distribution of natural and non-natural causes of death by age, 2023	32
Figure 4.4 - Percentage (%) of deaths due to communicable diseases (Group I), non-communicable diseases (Group II) and injuries (Group III) by year of death, 2002–2023*	35
Figure 4.5 - Percentage (%) of deaths due to communicable diseases (Group I), non-communicable diseases (Group II) and injuries (Group III) by sex and age group, 2023*	37
Figure 4.6 - Distribution of deaths for the leading causes of death by year of death and sex, 2021–2023*	41
Figure 4.7 - Distribution of other external causes of accidental injury, 2019–2023*	53

Abbreviations/acronyms

AIDS	Acquired Immuno Deficiency Syndrome
ANACoD	Analysing Mortality and Causes of Death
CDC	Centre for Disease Control
COVID-19	Corona Virus Disease of 2019
CRVS	Civil Registration and Vital Statistics
DHA	Department of Home Affairs
ECA	Economic Commission for Africa
GBD	Global Burden of Diseases
HIV	Human Immunodeficiency Virus
ICD-10	International Classification of Diseases 10 th Revision
ICD-11	International Classification of Diseases 11 th Revision
MACOD	Mortality and Causes of Death
MDR-TB	MultiDrug-Resistant Tuberculosis
MMDS	Mortality Medical Data System
NCDs	Non-Communicable Diseases
NDP	National Development Plan
NPR	National Population Register
Stats SA	Statistics South Africa
TB	Tuberculosis
UNECA	United Nation's Economic Commission for Africa
WHO	World Health Organization
XDR-TB	Extensively Drug-Resistant Tuberculosis

1. Introduction

1.1 Background

The responsibility of a well-established civil registration and vital statistics system (CRVS) is to attach a discrete legal identity to every individual in society (Peters, 2016). Over and above this basic function, CRVS provides essential data for policymaking, it is the foundation from which evidence-based decisions and policies can be formulated. Such a system provides a foundation for human rights and identity, data for development planning, monitoring of health and development goals, improving health systems, disaster response and humanitarian action (AbouZahr *et al.* 2017).

The CRVS system in South Africa is a well-established framework, with near-perfect registration completeness rates for both births and deaths. Its sustainability is supported by a strong legal framework that effectively captures both hospital and community deaths. The registration of deaths in South Africa is governed by the Births and Deaths Registration Act No. 51 of 1992 (Republic of South Africa, 1992). The Statistics Act No. 6 of 1999, as amended through Act 29 of 2024, gives Statistics South Africa (Stats SA) the legal mandate to publish vital statistics based on births and deaths reported through the civil registration system (Republic of South Africa, 2024).

Stats SA, in collaboration with DHA, National Department of Health (NDoH) and other stakeholders, has ensured continued provision of mortality data from the civil registration system. Sustained partnerships between the stakeholders are essential for improvements in mortality statistics to be realised and the findings from this statistical release will inform efforts aimed at strengthening civil registration and vital statistics systems in the country.

The utility of mortality data depends largely on the accuracy and completeness of cause-of-death reporting. The basis of which is the accurate completion of death notification forms by medical doctors, aligning with the International Classification of Diseases (ICD) for standardised reporting of causes of death which is gold standard for cause of death recording. Equally, timely information is critical for interventions and during health emergencies, it is key to informing health leaders on areas for intervention. Mortality and causes of death continue to be one of the key outputs for Stats SA. Consistent efforts to address the four-year backlog in the processing of MACOD are underway in order to meet its mandate to provide reliable information on the levels and causes of mortality through the application of appropriate quality criteria and standards, classifications and procedures for vital statistics (Republic of South Africa, 1999).

Accurate and reliable mortality data are also fundamental for understanding disease patterns, improving health outcomes, public health planning, and monitoring progress towards health and development goals. It is one of the indicators of the well-being and health status of a population, hence the South African government's flagship plan, National Development Plan (NDP), has "Health care for all by 2030" as one of the key development objectives (National Planning Commission, 2012).

1.2 Objectives of this statistical release

The mortality and causes of death statistical release is part of a regular series published by Stats SA, based on data collected through the civil registration system. This statistical release has two main objectives:

- To outline emerging trends spanning a 22-year period (2003–2024) and differentials in mortality by selected socio-demographic and geographic characteristics for deaths that occurred in 2024; and
- To present statistics on the causes of death for deaths that occurred in 2024, focusing on the underlying causes of death.

1.3 Scope of this statistical release

This release is based on information on mortality and causes of death from the South African civil registration system. All death notification forms from the DHA for deaths that occurred in 2024 or earlier that reached Stats SA during the 2025/2026 processing phase are covered. However, the main focus is on deaths that occurred in 2024. Deaths that occurred during the period 2003 to 2023 are also provided to show trends in mortality and causes of death. This release excludes stillbirths, which are also collected through the civil registration system using the same death notification form. The definitions of technical terms used in this release are provided in Appendix A.

1.4 Organisation and presentation of this statistical release

This release is composed of five sections. The first section consists of information on the background and purpose of the release. Section two lays out the data and methods, which focuses on data sources, including methods used in data processing, data editing, quality assurance and data analysis. The third section on registered deaths presents mortality levels, trends and differentials, specifically focusing on the socio-demographic and geographic characteristics of the deceased. The fourth section mainly covers information on the underlying causes of death for 2024 death occurrences. In addition, the section provides information on immediate, contributing, and underlying causes of death differentials by natural versus non-natural causes, as well as the Global Burden of Disease (GBD). Due to fewer proportions of COVID-19 deaths, there will be no dedicated sub-section for COVID-19, as was the case in 2020 and 2021. Finally, the last section presents a summary of the findings and concluding remarks.

This release is accompanied by additional analysis in Appendices D to Q under the tab “*Additional downloads*” on the Stats SA website. The workbook provided covers information on deaths and causes of deaths (including main group, broad group and underlying causes of death by population group), and the analysis is disaggregated by province and district municipality of death. Any information not provided in this release, or additional downloads may be requested from Stats SA. Enquiries may be made on any information not provided in this statistical release.

2. Data and methods

In this section, focus is placed on data sources, methods used in data processing, data editing and data analysis. Procedures followed in assessment of the quality of data are also covered.

2.1 Data source

This statistical release is based on administrative records from death notification forms acquired from the Department of Home Affairs (DHA). The DHA currently uses two death notification forms to register deaths: Form BI-1663 which was introduced in 1998 and Form DHA-1663 which was introduced in 2009 as a replacement of Form BI-1663. Form BI-1663 continues to be used in areas where it is still in stock. In the event that a medical practitioner could not certify the occurrence of death, a traditional leader (such as chief, induna, headman) completes Form DHA-1680 (referred to as the Death Report) to certify the occurrence of death and to provide a description of the circumstances that resulted in death. The completed Death Report is then sent to DHA where the information is transcribed on to the DHA-1663. During processing at Stats SA, the two death notifications (Form BI-1663 and Form DHA-1663) were merged into one dataset as data elements in these two forms are largely comparable. The main difference between the two forms is in the registration of perinatal deaths (stillbirths and deaths occurring within the first seven days of life). Form BI-1663 records perinatal deaths in the same section as all other deaths, whereas Form DHA-1663 has a separate section for the comprehensive recording of the details of perinatal deaths.

The Births and Deaths Registration Act, 1992 (Act No. 51 of 1992), amended in 2010 as the Births and Deaths Registration Amendment Act, 2010 (Act No. 18 of 2010) is the legislation governing the registration of deaths in South Africa (Republic of South Africa, 1992; Republic of South Africa, 2010). In addition, the 2014 Births and Deaths Regulations which repealed the 1992 Regulations prescribe that notice of death or stillbirth for all deaths that occur must be given within 72 hours of death occurrence by an informant, regardless of citizenship status of the deceased (Republic of South Africa, 1992). After death registration is completed, the DHA issues a death certificate to the informant and updates the National Population Register (NPR). The NPR only includes deaths for South African citizens and permanent residents whose birth records were already captured onto the NPR prior to death. Those not eligible for inclusion in the NPR are non-South African citizens who sojourn temporarily in the country and all South African citizens and permanent residents who died before notice of their births had been registered. Statistics South Africa (Stats SA) collects all death notification forms, irrespective of the deceased's citizenship status for processing, analysis and dissemination of mortality and causes of death information.

This statistical release is based on a total of 474 366 deaths that occurred in 2024 and 14 851 late death registrations for 2003 to 2023 that were registered at the DHA and reached Stats SA in time for the 2025/2026 processing phase.

2.2 Data processing

The processing of the completed death notification forms takes place at the Stats SA Data Processing Centre. The stages of data processing start with sorting of forms by year of death occurrence, pasting unique identifier labels on each form, data capturing; and coding socio-demographic and causes of death variables.

2.2.1 Classification of the causes of death

The cause-of-death statistics in this publication are compiled using the International Classification of Diseases (ICD), 10th Revision 2016 and 2019 Editions. The ICD is a system of categories to which morbid entities of either external or pathological causation are assigned according to established criteria. It is developed collaboratively between the World Health Organization (WHO) and various international agencies who are involved in mortality data quality improvement. It is revised from time to time in line with new adaptations, classifications and glossaries. All member states of the United Nations, including South Africa, agreed to use ICD as the standard classification system for compiling morbidity and mortality statistics. The South African National Information System also adopted it as a standard.

The primary purpose of ICD is to provide for conversion of word descriptions of diseases or conditions into an alphanumeric code, which permit easy storage, retrieval and analysis of data. It also allows for the systematic and standardised recording, analysis, interpretation, comparison and sharing of morbidity and mortality data within a population and across countries. The ICD-10 provides for coding and classification of diseases and injuries and a wide range of signs, symptoms and other abnormal findings (WHO, 2014a).

According to WHO (2016), the most effective public health objective is to prevent the underlying cause of death from operating. For this purpose, the WHO recommends that countries use the international form of medical certificate of cause of death to facilitate the selection of the underlying cause of death. The ICD-10 contains about 8 000 categories of causes of death, which are organised into 22 chapters consisting of communicable diseases, non-communicable diseases, ill-defined causes of death and external causes of injury and death (WHO, 2016).

Each chapter contains three-character categories which is subdivided into 10 four-character subcategories. However, for international comparisons, three-character coding is the mandatory level for reporting morbidity and mortality statistics, while four-character coding is recommended for more specific details about the disease or condition resulting in morbidity or mortality. Statistics South Africa codes the causes-of-death data at four-character level where sufficient details about the causes of death were available. However, this statistical release analyses up to three-character level.

The quality of causes of mortality statistics depends on completeness and accuracy of certified death notification forms. Coders at Stats SA follow the principle of 'what you see is what you code' when coding causes-of-death statistics. The coders use the ICD-10 for categories of causes of death coded in the ICD-10 manual. For categories that are not coded in the ICD-10 manual, Stats SA has outlined specific guidelines and

procedures. For example, according to these rules and procedures, immunosuppression is coded as immunodeficiency and not as human immunodeficiency virus (HIV) disease.

Medical practitioners sometimes report the cause of death as acquired immune suppression which is not coded in the ICD-10 manual. Based on the Stats SA guidelines, this is coded as human immunodeficiency virus (HIV) disease (B20-B24). Multidrug-resistant tuberculosis (MDR-TB) and extensively drug-resistant tuberculosis (XDR-TB) were assigned the ICD-10 special codes U51 and U52, respectively, and are included in the tuberculosis (A15-A19) broad group causes of mortality.

2.2.2 Generation of the underlying causes of death

The underlying cause of death is defined as: “(a) the disease or injury that initiated the sequence of events leading directly to death, or (b) the circumstances of the accident or violence that produced the fatal injury” (WHO, 2016: 31).

Stats SA uses IRIS software for the automated derivation of the underlying causes of death according to the ICD-10 rules. IRIS is the most up-to-date software available for coding as Mortality Medical Data System (MMDS) is no longer being supported by the US Centre for Disease Control who have moved to join the IRIS development core group. IRIS software contains language-dependent tables that can be developed to suit individual jurisdictions e.g. can include common phrases used in South African death certificates which may not be included in the standard English dictionary released with the IRIS package.

In instances where the software fails to derive the underlying cause of death, experienced coders at Stats SA derive the underlying cause of death manually within IRIS software. In efforts to improve the quality, timeliness, and consistency of coded data, IRIS was used to its full potential (i.e coding of the causes and derivation of the underlying cause of death).

2.3 Data editing

On completion of all data processing stages, the Stats SA data editing rules document was used to check for data consistency and accuracy and to flag implausible causes of death for further investigations. In addition, two electronic tools both developed by WHO: Analysing mortality levels and causes-of-death (ANACoD) version 3.0 and CoDEdit version 1.0 were used to further check data consistency and plausibility (WHO, 2014a and WHO, 2014b). The tools were developed to enhance the value of mortality statistics in informing health policies and programmes. Both tools were used to automatically check the 2023 death data for accuracy and consistency by highlighting cases with causes that were unlikely to cause death categorised by age and sex (sex-specific causes, age-specific causes and notifiable diseases) and possible misuse of ICD-10 codes as well as providing a summary of the records within the dataset (WHO, 2014a; WHO, 2014b). For example, for causes of death that are specific to one sex, the tools warn and flag for errors when the combination of sex and cause is wrong. The errors that were flagged by the tools were manually investigated (checked on the original death notification form) for verification and corrections were made where necessary.

The main difference between the two tools is that CoDEdit assesses data consistency and plausibility for each unit record while ANACoD checks the data at an aggregate level. However, this does not undermine the importance of ANACoD as it can calculate demographic indicators such as crude death rates, life expectancy and death registration completeness level.

2.4 Assessment of the quality of data

The importance of producing quality mortality statistics derived from the civil registration system cannot be over-emphasised, since they are the only source of health information data continuously available at national and local administrative levels.

The usability of mortality statistics wholly depends on their quality, while the data have potential to support decentralised population health administration, (WHO, 2013). An accurate, complete and timely civil registration system provides the foundation to produce reliable and routine vital statistics. However, the data can suffer from a range of quality limitations such as the extent of late registrations, timeliness of death registration, completeness of death registration, timeliness of publishing, accuracy of reporting, ill-defined causes of death and misreporting or misclassification of causes of death. It is therefore vitally important to check the data quality and to be transparent about data limitations, so that areas of improvement can be identified.

For the purpose of this statistical release, in addition to the ANACoD and CoDEdit electronic tools, the framework proposed by Mahapatra et al. (2007) was used to assess the quality of the 2023 causes of death data. A detailed assessment is provided in Appendix C.

2.5 Data analysis

A two-pronged approach was followed for this release, which includes mortality analysis and causes of death analysis. The first section on mortality describes information on selected socio-demographic variables and mortality patterns, based on frequency distributions and cross-tabulations.

The section further covers demographic indicators such as sex ratio at death, age-specific death rates and median ages at death. Sex ratios at death show the ratio of male deaths per 100 female deaths and age-specific death rates show variations in mortality taking into consideration the population size of each age group. Age-specific death rates indicate the number of deaths in a particular age group per 1 000 population in that age group while the median ages at death provide a basic measure of how early or late mortality occurs in a population over time.

The second section lays out analyses of information on causes of death, mainly based on ranking the natural underlying causes of death and proportions of deaths due to specific causes. The top-ranking causes determine the leading causes of death. The ranking indicates the frequency of causes of death among those causes eligible to be ranked and does not reflect causes of death in terms of their importance from a public health perspective. Causes of death with the same proportion received the same rank, and a rank was skipped

for the next cause. For example, if two causes of death had the same proportion and were ranked third, they both received the same rank, and the next cause received rank five.

The process of ranking natural underlying causes of death excluded symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00–R99), because such information is not sufficiently detailed to be of use for public health purposes. It is therefore essential to raise awareness among certifying practitioners to seek sufficient evidence to assign causes of these deaths to the more precise categories through training programmes and other initiatives. Due to concerns about violence and deaths due to accidents in South Africa, natural and non-natural causes have been separated. Although non-natural causes of death were not ranked, for analysis they were disaggregated by characteristics such as age, sex and province of death of the deceased that relay important information on the levels and patterns of non-natural deaths.

In addition, the second section also provides information on causes of death based on the Global Burden of Disease. Causes of deaths are categorised into three broad groups, namely Group I (communicable diseases), Group II (non-communicable diseases) and Group III (injuries). *Symptoms, signs and abnormal clinical and laboratory findings not elsewhere classified* (R00–R99) deaths which are ill-defined natural causes of death were accorded across communicable and non-communicable diseases categories.

3. Mortality

This chapter provides an analysis on the distribution of 2024 registered deaths that reached Stats SA during the 2025/2026 processing phase. The section mainly focuses on absolute numbers and percentage distributions of 2024 deaths by selected background characteristics of the deceased such as age, sex, place/institution of death and geographic information. Levels and trends of registered deaths over the period 2003–2023 are also included.


3.1 Levels and trends of mortality in 2024

Figure 3.1 shows that the total number of deaths that occurred and were registered at the DHA and processed by Stats SA for 2024 was 474 366, a decrease when compared with 483 298 deaths that were processed for 2023. This decline continues a downward trend observed after 2021, following a decrease in COVID-19-related deaths.

The general trend in the number of registered deaths processed by Stats SA indicates an increase from 2003 to 2006 when the number of deaths peaked at 614 864, and a decrease thereafter until 2019. A notable increase in the number of deaths was observed in 2020 (517 775), increasing to 624 528 deaths in 2021. This was followed by a sharp decrease in the number of deaths in 2022 (493 900), reaching pre-COVID-19 levels, and a further decline in 2024 (474 366).

The overall number of deaths per year increases as additional forms are processed at Stats SA. Additional forms may result from delayed registration or transmission of forms from DHA to Stats SA. It is, therefore, expected that additional forms, 2024 forms in particular, and for the previous years, will still be received for processing at Stats SA in the coming years. Updated information will be provided in the next statistical release.

Figure 3.1 - Number of registered deaths by year of death, 2003–2024*



*Data for 2003–2023 have been updated with late registrations / delayed death notification forms processed in 2025/2026

3.2 Age differentials

Table 3.1 below, shows the number and percentage distribution of deaths by age groups for deaths that occurred in 2024. The highest proportion of deaths were amongst those aged 65–69 years (9,1%), closely followed by those 70–74 years (8,9%) and 60–64 years (8,8%). The age groups which contributed the least number of deaths were age groups 5–9 years (0,5%) and 10–14 years (0,6%). Infants (age zero years) constituted 4,2% of the registered deaths.


Table 3.1 - Number and percentage (%) distribution of deaths by age group, 2024

Age group	Number	Percentage (%)
0	20 018	4,2
1–4	6 127	1,3
5–9	2 481	0,5
10–14	3 014	0,6
15–19	7 306	1,5
20–24	12 372	2,6
25–29	17 888	3,8
30–34	23 583	5,0
35–39	27 068	5,7
40–44	27 661	5,8
45–49	27 055	5,7
50–54	29 508	6,2
55–59	32 988	7,0
60–64	41 544	8,8
65–69	43 078	9,1
70–74	42 162	8,9
75–79	36 338	7,7
80–84	31 353	6,6
85–89	21 953	4,6
90+	20 869	4,4
Total	474 366	100,0

Figure 3.2 shows the percentage distribution of deaths by age groups and year of death between 2020 and 2024. A general observation is that the age pattern of mortality was somewhat consistent over the five-year period. However, slight deviations were observed during the COVID-19 years (2020 and 2021). During this time, the age groups between 0 years and 40–44 years recorded slightly lower proportions of deaths than typically observed. In contrast, the age groups between 55–59 years to 75–79 years showed a higher proportion of deaths than usual during the same period.

The overall pattern is generally characterised by high proportions of deaths amongst infants (zero years), lower proportions for ages 1–4 years, lowest proportions between 5–9 years and 10–14 years, rising but still low proportions between age group 15–19 years and 20–24 years. High proportions averaging over 6,0% are observed from age group 50–54 years to 80–84 years with decreasing proportions seen from age group 70 and older.

Figure 3.2 - Percentage (%) distribution of deaths by age group and year of death, 2020–2024*




*Excluding deaths with unspecified age
Data for 2020–2023 have been updated with late registrations / delayed death notification forms processed in 2025/2026

3.3 Sex differentials

Figure 3.3 shows the percentage distribution of deaths by sex and year from 2003 to 2024. Over the 22-year period, a consistent and persistent gender disparity in mortality is evident, with male deaths consistently exceeding those of females. In 2024, this pattern continued, with males accounting for 53,4% of deaths and females 46,6%. Notably, 2023 and 2024 recorded the widest gap in the series, marking the highest male share and the lowest female share of deaths observed.

The distribution of the deaths shows that there was a gap in the proportion of male and female deaths in 2003 (2,0 percentage points), narrowed yearly until 2005 (0,7 percentage points) and broadened again thereafter, reaching a male excess of 5,6 percentage points in 2016 to 2018. The emergence of COVID-19 in 2020 brought about the convergence in sex differentials, with the gap narrowing between the two sexes. The gap narrowed notably in 2020 (2,2 percentage points) and 2021 (2,0 percentage points), before widening in 2022 and 2023 to a high of 7,1 percentage points and remaining similarly high in 2024 (6,8 percentage points). Additional information on annual percentage changes in the number of deaths by sex, and Age-specific Death Rates (ASDRs) is provided in a workbook published separately, under additional downloads.

Figure 3.3 - Percentage (%) distributions of deaths by sex and year of death, 2003–2024*




*Excluding deaths with unspecified sex
Data for 2003–2023 have been updated with late registrations / delayed death notification forms processed in 2025/2026

3.4 Age and sex differentials

3.4.1 Distribution of deaths by age group and sex

Figure 3.4 shows the age and sex percentage distribution of deaths for 2024. It is observed that the proportion of deaths for males and females were both lowest and somewhat similar for the age groups 5–9 and 10–14 years. Overall, the male deaths exceeded those of females from age group 5–9 years up to 60–64 years, except for age group 65–69 years. From ages 70 years and above there were more female than male deaths. The gap in the proportion for male and female deaths was highest from 80 years and above, where female deaths surpassed male deaths by 3,9 percentage points at age group 80–84 years, by 3,8 percentage points at age groups 85–89 years and by 5,4 percentage points at ages 90 years and above. Additional information on absolute number of deaths by age, sex and year of death is provided in a workbook published separately, under additional downloads.

Figure 3.4 - Percentage (%) distribution of deaths by age group and sex, 2024*



*Excluding deaths with unspecified sex


3.4.2 Median ages at death by sex

The median age at death shows how early or late mortality occurs in a population and specifies the age at which half of the reported deaths occur. The median ages at death by sex and year of death over the years 2003 to 2024 are presented in Figure 3.5. Trends in median ages at death are important in the analysis of changes in mortality patterns over time, such as an increase in the proportion of deaths occurring at older ages or at younger ages. The former is indicative of improvements and postponement of mortality to older ages, while the latter indicates premature mortality.

Figure 3.5 shows that the median ages for males, females and total deaths decreased consistently from 2003 to 2005 and thereafter increased. It is also notable that both the declines and the increases were more pronounced amongst females relative to males. The median age for total deaths decreased from 43 years in 2003 to a low of 42 years in 2005, and then increased gradually from 2006, reaching a high of 60 years in 2021 before declining to 58 years in 2022. The median age increased again in 2024 to 60 years.

The median ages at death for males decreased from 44 years in 2003 to 43 years in 2005, while a reverse trend was observed between 2007 and 2021, from 44 years in 2007 to 57 years in 2021. In 2022, the median age at death for males decreased to 53 years and increased again to 55 years in 2024. Similarly, the female median ages at death decreased prior to 2006, from 42 years in 2003 to 41 years in 2005. Between 2006 and 2021, the average age at death for females increased from 43 years in 2006 to a high of 64 years in 2021. Similar to males, the female median age at death also dropped in 2022, recording 63 years and increased to 65 years in 2024.

Figure 3.5 - Median ages at death by sex and year of death, 2003–2024*



*Data for 2003–2023 have been updated with late registrations / delayed death notification forms processed in 2025/2026

3.4.3 Sex ratios by age groups


The sex ratio at death is an important demographic indicator, highlighting the number of male deaths relative to the number of female deaths. When there are equal numbers of male and female deaths, the sex ratio at death is equal to 100. If there are more male than female deaths, the sex ratio is above 100, and excess female deaths are indicated by a sex ratio at death less than 100.

Sex ratios at death by age group and year of death for the period 2020–2024 are shown in Figure 3.6. Over the five-year period, more male than female deaths were consistently observed from age 0 up to age group 65–69. Beyond this point, the pattern reversed, with more female deaths recorded than male deaths.

Results indicate that in 2023 and 2024, for the age groups 25–29 years to 35–39, the sex ratio showed a constant increase. In 2023 and 2024, the sex ratio at death reached a notable peak of 228 males per 100 females in the 20–24 age group, the highest across all age groups and years observed. A similarly high ratio of 220 was recorded in 2022 for the 15–19 age group.

Overall, from 2020 to 2024, the sex ratio generally showed an upward trend, particularly from age group 15–19 to age group 20–24. After the marked peaks observed at ages 15–19 in 2022 and 20–24 in 2023 and 2024, the sex ratio decreases steadily across the subsequent age groups, except for age group 50–54. This suggests that while male deaths remain higher than female deaths through much of the adult age range, the gap narrows progressively with increasing age. In the oldest age groups, the sex ratio falls below 100, indicating that female deaths exceed male deaths.

Figure 3.6 - Sex ratios by age group and year of death, 2020–2024*



*Excluding deaths with unspecified age and sex
Data for 2020–2023 have been updated to include late registrations/death notification forms processed in 2025/2026

3.5 Population group differences in mortality

Table 3.2 below, shows the distribution of the 2024 deaths by population group of the deceased. To account for the population composition, the table also shows deaths per thousand population. Black Africans had the highest percentage of deaths, comprising 65,1% of the total deaths, followed by the white population group (9,6%) and then the coloured population group (7,8%). The Indian/Asian population group (2,3%) accounted for the least percentage of registered deaths. The proportions observed are somewhat indicative of the variations in population size by population group.

While the percentage of deaths within the entire population indicates that the black African population group had the highest proportion of deaths, the death rate within each population group, reveals a different picture. The white population group had the highest death rate at 10 deaths per thousand population, followed by the coloured population group 7 deaths per thousand population and the black African and Indian/Asian population groups at 6,1 and 6,5 deaths per thousand respectively.

Additional information on the distribution of underlying causes of death by population group is provided in a workbook, which is published separately.

Table 3.2 - Number and percentage (%) distribution of deaths by population group, 2024

Population group	Number of deaths	Percentage (%) of deaths	Population group size	Deaths per thousand population
Black African	308 777	65,1	50 428 992	6,1
White	45 704	9,6	4 590 209	10,0
Indian/Asian	10 955	2,3	1 685 816	6,5
Coloured	37 087	7,8	5 278 488	7,0
Other	854	0,2		*
Unknown or unspecified*	70 989	15,0		*
Total	474 366	100,0	61 983 505	7,7

Source: 2024 Mid-year population estimates, 2026 series

*Deaths per thousand population not calculated for the "Unknown or unspecified" group due to the lack of a directly comparable population base

3.6 Marital status differences in mortality

Table 3.3 depicts the number and percentage distribution of deaths by marital status of the deceased. The majority of the deaths (38,3%) occurred amongst the people that had never been married, followed by those who are married (21,1%) and the widowed (10,7%). It is worth noting that the variations in the percentage of deaths by marital status may be affected by differences in population sizes across the marital status categories. In addition, caution should be exercised when interpreting the results, as 27,5% of the death notification forms had missing information on marital status of the deceased.

Table 3.3 - Number and percentage (%) distribution of deaths by marital status, 2024

Marital status	Number	Percentage (%)
Never married	181 539	38,3
Married	100 203	21,1
Widowed	50 666	10,7
Divorced	11 341	2,4
Unknown/unspecified/not applicable	130 617	27,5
Total	474 366	100,0

3.7 Differences in mortality by smoking status of the deceased

The number and percentage distribution of 2024 registered deaths classified by smoking status of the deceased is depicted in Table 3.4. below. Smoking status of the deceased is defined as the regular smoking of tobacco during the five years prior to death. The law strictly prohibits the sale of tobacco products to anyone under the age 16 years so the question on smoking status is applicable if the deceased was aged 16 years and older.

The table shows that the highest percentage of deaths were among people who were non-smokers (40,7%) while approximately 20,9% of the deaths occurred among people who were smoking. The table also shows that 32,2% of registered deaths in 2024 had unknown or unspecified smoking status. The high proportion of deaths with missing information on smoking status shows a poor reporting of this information on the death notification forms.

Table 3.4 - Number and percentage (%) distribution of deaths by smoking status among those aged 16 and older, 2024

Smoking status	Number	Percentage (%)
Yes	92 228	20,9
No	179 646	40,7
Do not know	27 479	6,2
Unknown or unspecified	142 304	32,2
Total	441 657	100,0

3.8 Differences in mortality by place or institution of death occurrence

The distribution of registered deaths by place or institution of death occurrence for 2024 is shown in Table 3.5 below. Results indicate that 40,6% of the deaths took place in hospitals, 2,2% occurred in emergency rooms or outpatient facilities, while 1,2% died in nursing homes. These three places of death occurrence account for 44,0% of total deaths that occurred within a health care facility. A total of 23,3% of all deaths occurred at home in 2024, while 2,1% were amongst people who had already died by the time they reached the hospital. A total of 28,8% of the death notification forms had unknown or unspecified information on place or institution of death.

Table 3.5 - Number and percentage (%) distribution of deaths by place of death occurrence, 2024

Place of death	Number	Percentage (%)
Hospital	192 745	40,6
Emergency room / Outpatient	10 205	2,2
Dead on arrival	10 155	2,1
Nursing home	5 660	1,2
Home	110 673	23,3
Other	8 318	1,8
Unknown/ unspecified	136 610	28,8
Total	474 366	100,0

3.9 Geographic variations in mortality

This section presents information on the distribution of registered deaths by province of occurrence and by the deceased's usual residence. The information was derived based on the 2020 municipal boundaries. Additional information on province of death by province of usual residence, sex and age group is provided in a workbook published separately.

3.9.1 Differences by province, age and sex

Table 3.6 shows the distribution of 2024 deaths by province of death occurrence and province of usual residence of the deceased. The province of death occurrence may not always be similar to the place of usual residence.

For province of death occurrence, the highest proportion of deaths (21,2%) occurred in Gauteng, followed by KwaZulu-Natal and Eastern Cape, each comprising 18,8% and 16,3%, respectively. The lowest percentage of deaths occurred in Northern Cape (3,1%).

With regard to province of usual residence, the same pattern was observed as that of province of death occurrence. Gauteng (20,3%) had the highest proportion of deaths, followed by KwaZulu-Natal (18,7%) and Eastern Cape (16,3%). The lowest percentage was Northern Cape at 3,0%.

It must be noted that analysis on geographic distribution of deaths is based only on place of death occurrence, not place of residence or place of birth of the deceased. However, information on the distribution of deaths by place of residence and place of birth of the deceased is available on request from Stats SA.

Table 3.6 - Distribution of deaths by province of death occurrence and province of usual residence of the deceased, 2024

Province	Province of death occurrence		Province of usual residence of deceased	
	Number	Percentage (%)	Number	Percentage (%)
Western Cape	57 757	12,2	56 011	11,8
Eastern Cape	77 315	16,3	77 180	16,3
Northern Cape	14 610	3,1	14 407	3,0
Free State	28 496	6,0	28 228	6,0
Kwa-Zulu Natal	89 002	18,8	88 827	18,7
North West	30 826	6,5	31 811	6,7
Gauteng	100 329	21,2	96 252	20,3
Mpumalanga	30 503	6,4	31 197	6,6
Limpopo	44 353	9,4	45 565	9,6
Outside South Africa	1 013	0,2	1 431	0,3
Unspecified	162	0,0	3 457	0,7
Total	474 366	100,0	474 366	100,0

4. Causes of death

This section presents information on causes of death for all deaths that occurred in 2024 and previous years and were registered in 2024, as well as comparisons with data from the previous years. In terms of the Birth and Deaths Act 51 of 1992, as amended, medical practitioners are responsible for certification of causes of death in the dedicated page for recoding the causes of deaths in the death notification form. Other role players are responsible for other sections of the form.

This section has nine sub-sections, namely: introduction, reported causes of death, method of ascertaining the cause of death, main groups of the underlying causes of death, natural and non-natural causes of death, major group of causes of death, broad groups of natural causes of death, non-natural causes of death, and comparison between immediate, contributing and underlying causes of death. A summary of causes of death by age, sex and province of death occurrence is also included in the analysis.

The 10th revision of the International Classification of Diseases (ICD-10) was used to classify the causes of death data in this publication. The analysis undertaken focuses mainly on underlying cause of death, which is defined as the disease or injury that initiated the train of events leading directly to death; or the circumstances of the accident or violence which produced the fatal injury (WHO, 1992). Previous publications have shown that Non-Communicable Diseases (NCD) pose a major barrier to health, quadrupling the burden of disease and as such, this necessitated the inclusion of analysis on the Global Burden of Diseases. Global Burden of Diseases is a critical resource for informed policymaking, as it provides a tool to quantify and compare the effects of different diseases in a population.

The last subsection presents comparison between the underlying, immediate and contributing causes of death. This analysis gives an overview of recorded instances of multiple causes of death and utilises information on all causes of death recorded on each death notification form, as death notification forms allow for reporting one or more causes of death on each form.

4.1 Reported causes of death

Table 4.1 shows information on the number of causes of death provided on each death notification form for deaths that occurred in 2024. Of the 474 366 forms 1 403 had no cause of death recorded on the forms. These are as a result of two scenarios:

Scenario 1: This is a situation where the page for recording causes of death information was missing when the form was received from the DHA, but there is a doctor's tick to show that it was a natural cause but with no specific cause given.

Scenario 2: This is where the page with causes of death information was missing when the form was received from the DHA, but the doctor indicated that the death was still under investigation and therefore cause of death had not yet been established.

In both scenarios, the age of the deceased was used to establish the underlying cause of death. ICD-10 codes such as R99 (other ill-defined and unspecified causes of mortality) and P96 (other conditions originating in the perinatal period) were used for these deaths, depending on the age of the deceased. ICD-10 code R99 (other ill-defined and unspecified causes of mortality) was used for adult deaths, while P96 (other conditions originating in the perinatal period) was used for perinatal deaths.

More than half of the death notification forms (52,6%) had only one cause recorded, followed by 23,9% of death notification forms which had two causes of death recorded. On 15,0% of the forms, three causes were recorded while 8,2% had four or more causes recorded.

Table 4.1 - Number and percentage (%) distribution of death notification forms by the number of causes reported on the death notification form, 2024

Number of the reported causes of death	Number of death notification forms	Percentage (%)
No cause	1 403	0,4
One cause	247 205	52,6
Two causes	114 500	23,9
Three causes	71 713	15,0
Four or more causes	39 545	8,2
Total	474 366	100,0

4.2 Method of ascertaining cause of death

The death notification form makes provision for a certifying official to indicate the method used to ascertain the cause of death. The eight options available on the form for method used to ascertain the death are shown in Table 4.2. In instances where the method of ascertainment was not filled but, on the form, there is date of post-mortem, the date was used as proxy for method of ascertainment in this case post-mortem.

The opinion of the attending medical practitioner was the most common method of ascertaining causes of death at 53,0%. It was followed by the opinion of the attending medical practitioner on duty at 14,3% and interview of family member at 11,9%. An autopsy was performed and results available in 12,3% of the 2024 deaths. There were 4,2% of forms indicated that cause of death was ascertained through postmortem examination, while 2,1% forms that had unspecified method of ascertainment.

Table 4.2 - Number and percentage (%) distribution of deaths by method used to ascertain the cause of death, 2024

Method of ascertaining the cause of death	Number	Percentage (%)
Autopsy	58 452	12,3
Post mortem examination	19 958	4,2
Opinion of attending medical practitioner	251 406	53,0
Opinion of attending medical practitioner on duty	68 066	14,3
Opinion of registered professional nurse	3 923	0,8
Interview of family member	56 449	11,9
Other	3 218	0,7
Autopsy results may be available later*	32	0,0
Autopsy not performed*	1 685	0,4
Unknown	1031	0,2
Unspecified	10146	2,1
Total	474 366	100,0

*For perinatal deaths only

4.3 Main groups of the underlying causes of death

This section presents an overview of the underlying causes of death for main groups (chapters) of classification of causes of death. The ICD-10 classifies diseases and related health problems into 22 chapters, of which 19 are used in the reporting of information on underlying causes of death (see Table 4.3).

This report excludes ICD-10 chapters 19, 21 and 22 and these are discussed briefly below:

- I. Chapter 19: *Injury, poisoning and certain other consequences of external causes (S00-T98)*. These codes are used to classify causes of death in other causes but not in the underlying causes.
- II. Chapter 21: *Factors influencing health status and contact with health services (Z00-Z99)*. These are only used in morbidity coding.
- III. Chapter 22: Codes for special purposes. These codes are used by WHO for the provisional assignment of new diseases of uncertain aetiology. U51 and U52 were used for coding *multidrug-resistant tuberculosis (MDR-TB)* and *extensively drug-resistant tuberculosis (XDR-TB)* in this release for individual causes of death but were both recoded to the broad group of tuberculosis (A15-A19) in the analyses.

Just as with other countries, South Africa (Stats SA) adopted the inclusion of COVID-19 as an additional chapter to the existing 22 chapters. This makes it easier for international comparison between countries and enables South Africa to align with countries using a similar approach. This is because the 2016 version of ICD-10 did not have COVID-19 as a chapter on the underlying causes of death.

The main groups (chapters) of the classification of causes of death are shown in Table 4.3 as both the number and percentage distribution of deaths. The most common main group of causes of death in 2024 was *symptoms and signs not elsewhere classified*, comprising 18,2% of all deaths. This main group consist mainly of information about various symptoms and signs that may not fit neatly into other categories; for example, some common conditions under this group include abnormal heart sounds (R01) and abnormal blood chemistry (R79).

The second most common main group of causes of death was *diseases of the circulatory system* (17,8%) followed by *certain infectious and parasitic diseases* (13,5%) of deaths. The fourth most common main group of causes of death was *external causes of morbidity and mortality* (13,3%). The rest of the groups contributed less than 10,0% of deaths in 2024. Amongst these were *neoplasms* (9,8%), *diseases of the respiratory system comprising* (8,2%) of all deaths, and *endocrine, nutritional and metabolic diseases* (7,4%), amongst others.


Table 4.3 - Distribution of deaths by main causes of death, 2024

No.	Main groups of underlying causes of death (based on ICD-10)	Number	Percentage (%)
18	Symptoms and signs not elsewhere classified (R00-R99)	86 477	18,2
9	Diseases of the circulatory system (I00-I99)	84 516	17,8
1	Certain infectious and parasitic diseases (A00-B99)	64 196	13,5
20	External causes of morbidity and mortality (V01-Y98)	62 871	13,3
2	Neoplasms (C00-D48)	46 264	9,8
10	Diseases of the respiratory system (J00-J99)	38 705	8,2
4	Endocrine, nutritional and metabolic diseases (E00-E90)	35 240	7,4
14	Diseases of the genitourinary system (N00-N99)	11 402	2,4
11	Diseases of the digestive system (K00-K93)	11 349	2,4
6	Diseases of the nervous system (G00-G99)	9 459	2,0
16	Certain conditions originating in the perinatal period (P00-P96)	8 828	1,9
3	Diseases of the blood and immune mechanism (D50-D89)	4 464	0,9
5	Mental and behavioural disorders (F00-F99)	3 849	0,8
17	Congenital malformations (Q00-Q99)	2 399	0,5
13	Diseases of the musculoskeletal system etc. (M00-M99)	1 904	0,4
12	Diseases of the skin and subcutaneous tissue (L00-L99)	1 426	0,3
15	Pregnancy, childbirth and puerperium (O00-O99)	518	0,1
	COVID-19 (U071-U072)	450	0,1
8	Diseases of the ear and mastoid process (H60-H95)	42	0,0
7	Diseases of the eye and adnexa (H00-H59)	7	0,0
	Total	474 366	100,0

*Including deaths due to *MDR-TB* and *XDR-TB*

A three-year (2022–2024) trend analysis on the distribution of deaths by main groups of causes of death is shown in Figure 4.1. It was observed that the rankings of the main groups of causes of death by year have remained more or less the same during the period 2022–2024. Throughout 2022–2024, the leading causes of death remained largely consistent, including *symptoms and signs not elsewhere classified*, *diseases of the circulatory system*, *certain infectious and parasitic diseases*, and *external causes of morbidity and mortality*. The most notable change occurred in 2022, when *diseases of the circulatory system* temporarily surpassed *symptoms and signs not elsewhere classified* as the top cause of death.

Less common causes, such as *certain conditions originating in the perinatal period*, *diseases of the blood and immune mechanism*, *diseases of the nervous system*, *diseases of the genitourinary system*, *diseases of the digestive system* and *congenital malformations*, consistently accounted for a small proportion of deaths, highlighting areas where mortality remains low.

Figure 4.1 - Percentage (%) distribution of deaths by main groups of causes of death, 2022–2024*

* Data for 2022–2023 have been updated with late registrations/delayed death notification forms processed in 2025/2026

** Including deaths due to *MDR-TB* and *XDR-TB*

4.4 Natural and non-natural causes of death

The ICD-10 codebook classifies all causes of death from chapters 1 to 18 of the ICD-10 as natural causes and chapter 20 (V01-Y98) as non-natural causes. This section discusses both natural and non-natural causes of death. Non-natural causes of death comprise all deaths that were not attributable or may not have been attributable to natural causes. Natural and non-natural causes of death information reported in this release was derived from the underlying causes of death based on specific causes of death recorded on the death notification form.

Table 4.4 shows that since 2003, the number of deaths due to natural causes were higher than the number of deaths due to non-natural causes. Between 2003 and 2006, there was a consistent increase in the number of natural deaths, after which a decline was observed until 2019 and increased again in 2020 and 2021 the decreased again in 2022–2024. The increase in natural deaths during 2020 and 2021 was attributed to COVID-19 deaths. Although the number of deaths due to non-natural causes fluctuated over the years, the overall trend from the start to the end points shows an increase, rising from 52 992 in 2003 to 62 871 in 2024.


Table 4.4 - Number of natural and non-natural deaths by year of death occurrence, 2003–2024*

Year of death	Number of natural deaths	Number of non-natural deaths	Total
2003	506 247	52 992	559 239
2004	525 696	53 484	579 180
2005	546 211	54 095	600 306
2006	561 537	53 327	614 864
2007	552 211	54 671	606 882
2008	545 216	53 716	598 932
2009	533 808	50 910	584 718
2010	503 982	49 498	553 480
2011	473 391	47 616	521 007
2012	449 268	49 276	498 544
2013	430 372	50 243	480 615
2014	429 074	51 364	480 438
2015	424 778	53 896	478 674
2016	421 289	54 263	475 552
2017	415 792	54 801	470 593
2018	415 363	56 884	472 247
2019	410 275	58 549	468 824
2020	467 811	49 970	517 781
2021	562 907	61 624	624 531
2022	427 789	66 129	493 918
2023	418 704	64 740	483 444
2024	411 495	62 871	474 366

*Data for 2003–2023 have been updated with late registrations/delayed death notification forms processed in 2025/2026

Figure 4,2 reaffirms that through the entire 2003 to 2024 period, the proportion of deaths due to natural causes have always been higher than that of non-natural causes. The pattern relatively remained the same throughout this period with the exception of the years where COVID-19 was at its heights. These were 2020 and 2021 where there was a significant decrease in the proportion of deaths due to non-natural causes and this was mainly driven by COVID-19 deaths which caused a significant increase in the number of natural deaths.

Figure 4.2 - Percentage (%) distribution of natural and non-natural causes of death by year of death, 2002–2024*




*Data for 2003–2023 have been updated with late registrations/delayed death notification forms processed in 2025/2026

4.4.1 Natural and non-natural causes of death by age

The percentage distribution of deaths due to natural and non-natural causes classified by age group for deaths that occurred in 2024 is provided in Figure 4.3. The general pattern observed is that the proportion of deaths due to non-natural causes generally increased consistently from age 0 (3,6%) to age group 20–24 (50,2%) then decreased thereafter. It is worth noting that age group 20–24 was the age mostly affected by non-natural causes, with over half of the deaths in this age group due to non-natural causes. Other ages with higher proportions (over 30,0%) of deaths due to non-natural causes were age groups 25–29 (47,9%), 15–19 (44,0%), 30–34 (39,0%), 5–9 (35,7%) and 10–14 (34,4%). Ages least affected by non-natural deaths were infants (less than 0) and older ages (55 years and older) where less than 10,0% of the deaths in each of these age groups were due to non-natural causes of death.

Natural deaths are very high in infancy, accounting for 96,4% of deaths. From age groups 1–4 to 20–24 deaths due to natural causes steadily decline, with the 20–24 age group showing nearly equal proportions of natural (49,8%) and non-natural (50,2%) deaths. Natural deaths generally increase from age group 25–29 (52,1%) up to 70–74 (97,6%), after which they remain relatively stable at around 97,0%.

Figure 4.3 - Percentage (%) distribution of natural and non-natural causes of death by age, 2024



4.5 Major groups of causes of death as per Global Burden of Disease

The Global Burden of Disease (GBD) study is an all-inclusive program of disease burden that assesses mortality and disability from major diseases, injuries, and risk factors. It provides a comprehensive picture of mortality and disability across countries, time, age, and sex and is a landmark initiative that systematically quantifies the prevalence, morbidity, and mortality for hundreds of diseases, injuries, and risk factors of global health importance (Institute for Health Metrics and Evaluation). This is a useful measure as countries can combine this type of evidence along with information about policies and their costs to decide how to set their health targets and interventions. GBD also makes comparisons across populations, enabling understanding of the changing health challenges facing people across the world.

The twenty-two ICD-10 chapters used in the reporting of information on underlying causes of death can be further condensed into three groups of causes of death as per the Global Burden of Disease cause list:

Group I:

- Communicable diseases (e.g. *tuberculosis, pneumonia, diarrhoea, malaria, measles*);
- Maternal and perinatal causes (e.g. *maternal hemorrhage, birth trauma*); and
- Nutritional conditions (e.g. *protein-energy malnutrition*).

Group II: Non-communicable diseases (e.g. *cancer, diabetes, heart disease and asthma*)

Group III: External causes of mortality (e.g. *accidents, homicide and suicide*)


Communicable diseases are diseases caused by pathogenic microorganisms, such as bacteria, viruses, parasites or fungi and can be spread, directly or indirectly, from one person to another. These include, amongst other diseases, diarrhoea, tuberculosis and pneumonia. Non-communicable diseases are medical conditions or diseases that are non-infectious or non-transmissible among people. These diseases last for longer periods of time and progress slowly and include, amongst others, cancer, asthma and heart diseases. External causes of mortality are the non-natural causes of death which are discussed in chapter 20 of the ICD-10.

The percentage distribution of deaths by group type and year of death are depicted in Figure 4.4. The pattern shows that in 2003; there was no significant difference in the proportion of deaths due to communicable and non-communicable diseases. Starting from the year 2004 up to 2008, deaths due to communicable diseases surpassed non-communicable deaths with a percentage point difference ranging between 2,8 and 5,2. In 2009, the proportion of deaths due to communicable and non-communicable diseases were almost exactly the same in terms of proportion. From 2010 to 2018, the gap between the communicable and non-communicable diseases became wider with more deaths resulting from non-communicable diseases, with a percentage point difference ranging from 2,2 to 29,8. This however changed in the years 2019 to 2021, a convergence was observed between communicable and non-communicable diseases. This was mainly driven by the high number of COVID-19 deaths especially for the years 2020 and 2021.

From 2022, the pattern reversed back to the one observed prior to COVID-19 years where the gap between communicable and non-communicable diseases widened, reaching a high of 34,7 percentage point difference between communicable and non-communicable diseases in 2024.

The figure also shows that non natural causes of death decreased between 2003 and 2006. The increase in the proportion of non-natural deaths was again consistently observed from 2010 (8,9%) to 2019 (12,5%). This pattern was disturbed in 2020 and 2021, after which, proportions surpassing those of pre-COVID-19 were observed in 2023 and 2024, where non-natural deaths accounted for 13,4% and 13,3% of deaths respectively.

Figure 4.4 - Percentage (%) of deaths due to communicable diseases (Group I), non-communicable diseases (Group II) and injuries (Group III) by year of death, 2003–2024*




*Data for 2003–2023 have been updated with late registrations/delayed death notification forms processed in 2025/2026

The percentage distribution of group type by sex and age group is presented in Figure 4.5. Infants (aged 0 years) had the highest proportion of deaths due to communicable diseases for both sexes (76,0% for males and 74,5% for females). Deaths due to communicable diseases peak again at ages 40–44 (35,9%) for males while it peaks again from ages 30–34 (48,8%) for females. The proportion of deaths due to communicable diseases declines gradually with age from the age of 45 for males and from the age of 35 for females. Communicable diseases were lowest amongst the elderly at age group 80–84 for males (13,9%) and at age group 75–79 (12,6%) for females.

The trend in the proportion of deaths due to non-communicable diseases increases from age groups 30–34 for males (21,6%) and 25–29 years for females (31,5%), until older ages.

The most notable pattern in non-natural causes of death i.e. external causes of death including accidents and violence, for both sexes is the steady increase and peak at age group 20–24 years for males (61,9%), and age group 5–9 years amongst females (30,9%). Deaths due to external causes were much higher for males than for females starting from age group 0 and through all age groups. For males, non-natural causes of death declined to below 10,0% from age group 60–64 years, while for females the decline is observed from age group 40–44 years.

Figure 4.5 - Percentage (%) of deaths due to communicable diseases (Group I), non-communicable diseases (Group II) and injuries (Group III) by sex and age group, 2024*



4.6 Broad groups of natural causes of death

This subsection presents information on the leading underlying natural causes of death for broad groups using top ten ranking. The ten leading causes were identified by ranking the causes of death by the number of deaths among those eligible for ranking as described in Section 2 and excludes symptoms, signs and abnormal findings, not elsewhere classified as well as all non-natural deaths (external causes of morbidity and mortality). The top-ranking causes determine the leading underlying natural causes of death as it accounts for large numbers of deaths within a specified population and time period.

Ranking causes of death is crucial for identifying major health challenges and allocating resources for prevention and treatment. Unspecified causes don't offer actionable insights for these purposes. Knowing that a large number of deaths are attributed to "unspecified" conditions doesn't help in designing targeted interventions. By focusing on specific, well-defined causes, public health officials can better track trends, identify risk factors, and evaluate the effectiveness of health programs.

4.6.1 Overall pattern of the leading underlying natural causes of death

The ten leading underlying natural causes of death in South Africa for the years 2022–2024 are shown in Table 4.5 below. The previous years 2022 and 2023 have been included in the analysis to show recent trends in natural causes of death. The table provides trends in the ten leading underlying causes of death by absolute numbers and percentages over the three-year period.

Table 4.5 shows that nine of the ten leading causes of death in 2024 were the same for the three-year period, although they differed in rank as well as proportions. The COVID-19 pandemic first reached South Africa in 2020 and became the leading cause of death in 2020 was completely out of the top ten leading underlying causes of death for the years 2002 to 2004. This gives a reflection that acute phase of the COVID-19 pandemic is now over, however, it is still not completely eradicated as there were still significant number of deaths due to COVID-19 in 2022 and 2023 though they were not in the top ten.

In 2024, *diabetes mellitus* remained the top leading underlying cause of death, just as it was in the previous two years, responsible for 6,1% deaths in the country. It remained the leading underlying cause of death in the three-year period. The second leading underlying cause of death in 2024 was *hypertensive diseases* responsible for 5,5% deaths. In terms of rankings between the years 2022 and 2024, there were no notable changes between the three years. All the causes changed only 1 rank in between the years. For the years 2023 and 2024, six of the ten leading underlying causes of death kept the same rank and the other four changed, either going down or up the rank.

Table 4.5 – The ten leading underlying natural causes of death, 2022–2024*

Causes of death (ICD-10)	2022			2023			2024		
	Rank	Number	%	Rank	Number	%	Rank	Number	%
Diabetes mellitus (E10-E14)	1	33 126	6,7	1	27 967	5,8	1	28 767	6,1
Hypertensive diseases (I10-I15)	2	31 502	6,4	3	25 286	5,2	2	25 972	5,5
Cerebrovascular diseases (I60-I69)	3	29 084	5,9	2	25 844	5,3	3	25 539	5,4
Tuberculosis (A15-A19)	5	20 715	4,2	4	20 472	4,2	4	18 666	3,9
Human immunodeficiency virus [HIV] disease (B20-B24)	4	21 109	4,3	5	19 406	4,0	5	18 025	3,8
Influenza and pneumonia (J09-J18)	7	19 913	4,0	6	14 710	3,0	6	14 959	3,2
Other forms of heart disease (I30-I52)	6	20 513	4,2	7	14 211	2,9	7	14 059	3,0
Ischaemic heart diseases (I20-I25)	9	13 268	2,7	8	13 547	2,8	8	13 351	2,8
Chronic lower respiratory diseases (J40-J47)	10	11 993	2,4	10	12 090	2,5	9	12 081	2,5
Other viral diseases (B25-B34)	8	13 296	2,7	9	13 375	2,8	10	11 945	2,5
Other Natural		213 252	43,2		231 650	47,9		228 131	48,1
Non-natural		66 129	13,4		64 740	13,4		62 871	13,3
All causes		493 900	100,0		483 298	100,0		474 366	100,0

*Data from 2022–2023 have been updated with late registrations/delayed death notification forms processed in 2025/2026

**Including deaths due to MDR-TB and XDR-TB

... Category not in top ten

4.6.2 Leading underlying natural causes of death by sex

Table 4.6 shows the distribution of the ten leading underlying natural causes of death in 2024 by sex. Mainly, nine of the ten leading causes of death were the same for both sexes. The two underlying causes that were not common between the two sexes were *chronic lower respiratory diseases* and *malignant neoplasm of female genital organs*. *Chronic lower respiratory diseases* ranked eighth amongst males responsible for 3,0% male deaths while *malignant neoplasm of female genital organs* also ranked eighth amongst females responsible for 2,9% deaths. Among the top ten causes of death, *influenza and pneumonia* and *other viral diseases* were the only causes that had the same rankings for both sexes. *Influenza and pneumonia* ranked sixth, accounting for 3,2% of male deaths and 3,1% of female deaths, while *other viral diseases* ranked tenth for both sexes, accounting for 2,5% of deaths in both males and females.

The leading underlying cause of death for males was *tuberculosis* accounting for 4,8% male deaths followed closely by *cerebrovascular diseases* and *diabetes mellitus*, both accounting for 4,5% of male deaths. Causes that had the same proportion received the same rank and a rank was skipped for the next cause. For example, if two causes of death had the same proportion and were the top-ranking causes, they both received rank one, and the next cause received rank three.

The leading underlying cause of death for females was *diabetes mellitus* responsible for 7,9 % female deaths, followed closely by *hypertensive diseases* responsible for 7,1% female deaths. *Cerebrovascular diseases* ranked third responsible for 6,4% female deaths. For both sexes, six of the ten leading underlying causes of death are non-communicable diseases.

Table 4.6 - The ten leading underlying causes of death for males and females, 2024*

Causes of death (ICD-10)	Male			Female		
	Rank	Number	(%)	Rank	Number	(%)
Diabetes mellitus (E10-E14)	3	11 334	4,5	1	17 429	7,9
Hypertensive diseases (I10-I15)	4	10 322	4,1	2	15 644	7,1
Cerebrovascular diseases (I60-I69)	2	11 352	4,5	3	14 182	6,4
Human immunodeficiency virus [HIV] disease (B20-B24)	5	9 295	3,7	4	8 726	4,0
Other forms of heart disease (I30-I52)	9	6 624	2,6	5	7 428	3,4
Influenza and pneumonia (J09-J18)	6	7 998	3,2	6	6 934	3,1
Tuberculosis (A15-A19)	1	12 175	4,8	7	6 473	2,9
Malignant neoplasms of female genital organs (C51-C58)	...	...	...	8	6 295	2,9
Ischaemic heart diseases (I20-I25)	7	7 522	3,0	9	5 824	2,6
Other viral diseases (B25-B34)	10	6 336	2,5	10	5 607	2,5
Chronic lower respiratory diseases (J40-J47)	8	7 479	3,0	...	...	...
Other Natural		113 004	44,7		113 145	51,3
Non-natural		49 611	19,6		12 959	5,9
Total		253 052	100,0		220 646	100,0


* Excluding deaths with unspecified sex

**Including deaths due to *MDR-TB* and *XDR-TB*

...Category not in top ten

Figure 4.6 shows the percentage distribution of deaths associated with the ten leading causes of death classified by sex for the period 2022–2024. In 2024, *Tuberculosis* was the leading cause of death among males at 4,8%, followed by *cerebrovascular diseases* and *diabetes mellitus*, both at 4,5%. Among males, there were no causes of death that moved outside the top ten leading underlying causes of death in the 3-year period though their proportions differed between the years.

In 2024 the leading causes of death among females were dominated by chronic conditions. *Diabetes mellitus* was the leading cause of death and accounted for 7,9%, while *hypertensive diseases* came second and stood at 7,1%. *Cerebrovascular diseases* contributed 6,4%, reflecting a gradual decline from earlier years. Infectious diseases such as *HIV diseases* (4,0%) and tuberculosis (2,9%) remained significant but showed steady decreases. Meanwhile, malignant neoplasms of female genital organs continued an upward trajectory rising from 2,4% in 2022 to 2,9% in 2024.

Figure 4.6 - Distribution of deaths for the leading causes of death by year of death and sex, 2022–2024*

*(1) Data for 2022–2023 have been updated with late registrations/delayed death notification forms processed in 2025/2026

(2) Including deaths due to *MDR-TB* and *XDR-TB*

4.6.3 Leading underlying natural causes of death by age

Analysis of the broad age groups (0, 1–14, 15–44, 45–64, and 65 years and older) is recommended by the World Health Organization (WHO) for classifying ages for international comparison (WHO, 2009). Table 4.7 shows the ten leading underlying natural causes of death for these age groups. Further decomposition of age and leading underlying natural causes of death are provided in Tables 4.8 (under 5 years) and 4.9 (15–24 years).

Table 4.7 shows that only *influenza and pneumonia* was the only common underlying cause of death in all the age groups, although the ranking varied greatly by age. It was the leading cause of death for age group 1–14 responsible for 7,9% deaths in this age group, it ranked fourth for age group 15–44 accounting for 4,3%; it was ranked sixth for infants and seventh for those aged 65 years and older.

Respiratory and cardiovascular disorders specific to the perinatal period was the leading underlying cause of death for infant deaths (age 0) accounting for 15,5% infant deaths. The second leading underlying cause of death for infants was *other disorders originating in the perinatal period* responsible for 9,2% infant deaths. *Intestinal infectious diseases* ranked third, responsible for 6,2% infant deaths and ranked second for age group 1–14 responsible for 6,5% deaths in this age group.

The leading underlying cause of death for the age group 15–44 years was *human immunodeficiency virus [HIV] diseases*, constituting 12% of deaths, followed by *tuberculosis*, accounting for 10,8% of deaths. *Other viral diseases* which ranked third accounting for 7,3%. *Malignant neoplasms of female genital organs* which ranked ninth was in the top ten leading underlying causes of death only in this age group responsible for 1,7% deaths.

Seven of the ten leading causes of death for those 45–64 and 65 years and older were the same, with differences in rank and the contribution of each cause to the overall number of deaths in each age group. While *diabetes mellitus* was the leading cause of death among those aged 45–64, contributing 7,9% of deaths in this age group, it was the second leading cause of death among those aged 65 and older, accounting for 9,3% of deaths. Conversely, *hypertensive diseases* which was the leading cause of death for those aged 65 and older (9,6%), was the fifth leading underlying cause of death for those aged 45–64 (5,3%).

The underlying causes of death in age group 65 years and older that were not in the age group 45–64 are *other bacterial diseases*, *other forms of heart diseases* and *renal failure*. *Tuberculosis*, *Human immunodeficiency virus [HIV] disease* and *other viral diseases* were among the ten leading causes of death for the age group 45–64 age group, which were not listed for the age group 65 years and older.

Table 4.7 - The ten leading underlying natural causes of death for broad age groups, 2024

Causes of death (ICD-10)	0			1-14			15-44			45-64			65+		
	Rank	Number	%	Rank	Number	%	Rank	Number	%	Rank	Number	%	Rank	Number	%
Respiratory and cardiovascular disorders specific to the perinatal period (P20-P29)	1	2 985	15,5		...	...		...	...		...	...		...	...
Other disorders originating in the perinatal period (P90-P96)	2	1 784	9,2		...	...		...	...		...	...		...	...
Intestinal infectious diseases (A00-A09)	3	1 192	6,2	2	525	6,5		...	...		...	...		...	...
Infections specific to the perinatal period (P35-P39)	4	1 179	6,1		...	...		...	...		...	...		...	...
Disorders related to length of gestation and fetal growth (P05-P08)	5	1 015	5,3		...	...		...	...		...	...		...	...
Influenza and pneumonia (J09-J18)	6	958	5,0	1	636	7,9	4	3 204	4,3	10	3 691	3,1	7	6 470	3,4
Fetus and newborn affected by maternal factors and by complications of pregnancy, labour and delivery (P00-P04)	7	799	4,1		...	...		...	...		...	...		...	...
Congenital malformations of the circulatory system (Q20-Q28)	8	516	2,7		...	...		...	...		...	...		...	...
Haemorrhagic and haematological disorders of fetus and newborn (P50-P61)	9	437	2,3		...	...		...	...		...	...		...	...
Digestive system disorders of fetus and newborn (P75-P78)	10	403	2,1		...	...		...	...		...	...		...	...
Cerebral palsy and other paralytic syndromes (G80-G83)		...	...	3	317	3,9		...	...		...	...		...	...
Tuberculosis (A15-A19)		...	...	4	281	3,5	2	8 066	10,8	4	6 788	5,7		...	...
Malnutrition (E40-E46)		...	...	5	226	2,8		...	...		...	...		...	...
Episodic and paroxysmal disorders (G40-G47)		...	...	6	208	2,6	8	1 474	2,0		...	...		...	...
Other bacterial diseases (A30-A49)		...	...	7	189	2,3		...	...		...	...	10	3 574	1,9
Other forms of heart disease (I30-I52)		...	...	8	188	2,3	6	2 014	2,7		...	...	4	8 130	4,3
Human immunodeficiency virus [HIV] disease (B20-B24)		...	...	9	170	2,1	1	8 944	12,0	2	7 218	6,1		...	...
Inflammatory diseases of the central nervous system (G00-G09)		...	...	10	159	2,0		...	...		...	...		...	...
Other viral diseases (B25-B34)		...	...		...	...	3	5 419	7,3	6	4 949	4,2		...	...
Cerebrovascular diseases (I60-I69)		...	...		...	...	5	2 045	2,7	3	7 170	6,0	3	16 194	8,5
Diabetes mellitus (E10-E14)		...	...		...	...	7	1 616	2,2	1	9 364	7,9	2	17 728	9,3
Malignant neoplasms of female genital organs (C51-C58)		...	...		...	...	9	1 301	1,7		...	...		...	...
Renal failure (N17-N19)		...	...		...	...	10	1 288	1,7		...	...	9	5 062	2,7
Hypertensive diseases (I10-I15)		...	...		...	...		...	...	5	6 287	5,3	1	18 332	9,6
Ischaemic heart diseases (I20-I25)		...	...		...	...		...	...	7	4 188	3,5	5	7 921	4,2
Malignant neoplasms of digestive organs (C15-C26)		...	...		...	...		...	...	8	4 095	3,4	8	5 312	2,8
Chronic lower respiratory diseases (J40-J47)		...	...		...	...		...	...	9	3 750	3,1	6	7 200	3,8
Other Natural		8 022	41,6		5 196	64,2		39 331	52,7		61 694	51,8	50	94 291	49,6
All natural causes		19 290	100,0		8 095	100,0		74 702	100,0		119 194	100,0		190 214	100,0

*Including deaths due to *MDR-TB* and *XDR-TB*

... Category not in top ten

4.6.4 Leading underlying natural causes of death for children aged below five years by age groups

Ten leading causes of death for neonatal deaths (infants that died within the first 28 days of life), post-neonatal deaths (29 days to 11 months), all infant deaths (aged less than one year), and deaths among those aged 1–4 years are shown in Table 4.8. Infant deaths are composed of both neonatal and post-neonatal deaths.

Table 4.8 shows that neonatal deaths mainly resulted from *respiratory and cardiovascular disorders specific to the perinatal period*, which was the leading underlying cause of death among neonates, and responsible for 28,0% deaths. The second leading underlying cause of death for neonatal deaths was *other disorders originating in the perinatal period* accounting for 18,5% of all neonatal deaths, followed by *infections specific to the perinatal period*, responsible for 11,6% of deaths in this age group.

The leading cause of death for those who died during the post-neonatal period was *intestinal infectious diseases* (10,7%) followed closely by *influenza and pneumonia* (8,9%). These two causes were the highest contributors of post-neonatal deaths, accounting for 19,6% of deaths occurring during this period. *Respiratory and cardiovascular disorders specific to the perinatal period* (3,5%) was the third leading cause of death, and *malnutrition* was fourth responsible for 3,2% deaths in this age group.

Overall, for infants (less than one year), the leading underlying cause of deaths was *respiratory and cardiovascular disorders specific to the perinatal period* (14,9%). *Other disorders originating in the perinatal period* (8,9%) was ranked second. *Intestinal infectious diseases* (6,0%) and *infections specific to perinatal period* (5,9%) and by *disorders related to length of gestation and fetal growth* (5,1%) were ranked third, fourth and fifth, respectively.

The three leading causes of death for those aged 1–4 years were *influenza and pneumonia* (6,7%), *intestinal infectious diseases* (6,5%) and *malnutrition* (3,4%) ranking first, second and third, respectively. *Tuberculosis* (2,2%) was the fourth leading underlying cause of death while *metabolic disorders* (1,7%) was the fifth leading underlying cause of death.

For under five ages, *respiratory and cardiovascular disorders specific to the perinatal period* was the leading underlying cause of death responsible for 11,4% of deaths, followed by *other disorders originating in the perinatal period* (6,8%) while *intestinal infectious diseases* accounted for 6,1% of deaths in this age group. *Influenza and pneumonia* was ranked the fourth leading underlying cause of death amongst the under-5 mortality, responsible for 5,2% of deaths.

Table 4.8 - The ten underlying natural causes of death for infants and children aged below five years, 2024

Causes of death (ICD-10)	Neonatal (0-28 days)			Post-neonatal (29 days to 11 months)			Less than 1 year			1-4 years			Under 5 years		
	Rank	Number	%	Rank	Number	%	Rank	Number	%	Rank	Number	%	Rank	Number	%
Respiratory and cardiovascular disorders specific to the perinatal period (P20-P29)	1	2 609	28,0	3	376	3,5	1	2 985	14,9	...	...	...	1	2 990	11,4
Other disorders originating in the perinatal period (P90-P96)	2	1 730	18,5	...	...	...	2	1 784	8,9	...	...	...	2	1 789	6,8
Infections specific to the perinatal period (P35-P39)	3	1 086	11,6	...	...	...	4	1 179	5,9	...	...	...	5	1 180	4,5
Disorders related to length of gestation and fetal growth (P05-P08)	4	914	9,8	...	...	...	5	1 015	5,1	...	...	...	6	1 020	3,9
Fetus and newborn affected by maternal factors and by complications of pregnancy, labour and delivery (P00-P04)	5	794	8,5	...	...	...	7	799	4,0	...	...	...	7	800	3,1
Haemorrhagic and haematological disorders of fetus and newborn (P50-P61)	6	406	4,4	...	...	...	9	437	2,2	...	...	...	...	...	...
Other congenital malformations (Q80-Q89)	7	309	3,3	...	...	...	...	...	...	...	...	...	...	...	...
Digestive system disorders of fetus and newborn (P75-P78)	8	293	3,1	...	...	...	10	403	2,0	...	...	...	...	...	...
Congenital malformations of the circulatory system (Q20-Q28)	9	222	2,4	8	294	2,8	8	516	2,6	...	...	...	8	597	2,3
Chromosomal abnormalities, not elsewhere classified (Q90-Q99)	10	154	1,7	10	181	1,7	...	...	...	...	...	...	...	...	...
Intestinal infectious diseases (A00-A09)	...	...	...	1	1 143	10,7	3	1 192	6,0	2	401	6,5	3	1 593	6,1
Influenza and pneumonia (J09-J18)	...	...	...	2	948	8,9	6	958	4,8	1	408	6,7	4	1 366	5,2
Malnutrition (E40-E46)	...	...	...	4	342	3,2	...	...	...	3	210	3,4	9	559	2,1
Other bacterial diseases (A30-A49)	...	...	...	5	337	3,2	...	...	...	6	99	1,6	...	...	...
Other acute lower respiratory infections (J20-J22)	...	...	...	6	324	3,0	...	...	...	7	98	1,6	10	446	1,7
Metabolic disorders (E70-E90)	...	...	...	7	302	2,8	...	...	...	5	104	1,7	...	...	...
Other diseases of the respiratory system (J95-J99)	...	...	...	9	217	2,0	...	...	...	...	...	...	...	...	...
Tuberculosis (A15-A19)	...	...	...	...	...	...	...	...	...	4	134	2,2	...	...	...
Cerebral palsy and other paralytic syndromes (G80-G83)	...	...	...	...	...	...	...	...	...	8	94	1,5	...	...	...
Episodic and paroxysmal disorders (G40-G47)	...	...	...	...	...	...	...	...	...	9	89	1,5	...	...	...
Other forms of heart disease (I30-I52)	...	...	...	...	...	...	...	...	...	10	87	1,4	...	...	...
Other Natural		727	7,8		5 582	52,2		8 022	40,1		2 799	45,7		11 473	43,9
Non-natural		85	0,9		643	6,0		728	3,6		1 604	26,2		2 332	8,9
All causes		9 329	100,0		10 689	100,0		20 018	100,0		6 127	100,0		26 145	100,0

* Including deaths due to *MDR-TB* and *XDR-TB*

... Category not in top ten

4.6.5 Leading underlying natural causes of death for the population aged 15–24 years

According to the WHO recommendations, the 15–24 age group must also be included in the analysis for international comparison (WHO, 1992). This analysis is provided in Table 4.9. In 2024, *tuberculosis* was the leading cause of death for those aged 15–24, accounting for 5,0% of deaths, followed by *human immunodeficiency virus [HIV] disease* (3,4%) and *Influenza and pneumonia* (3,3%). *Other viral diseases* (2,3%), *episodic and paroxysmal disorders* (1,9%) *other forms of heart disease* (1,3%) and were the fourth, fifth and sixth leading causes of death, respectively. The ten leading causes of death in this age group contributed just over a quarter (21,2%) of deaths in this age group.

Table 4.9 - The ten leading underlying causes of death for the population aged 15–24 years, 2024

Causes of death (ICD-10)	15-24		
	Rank	Number	Percentage (%)
Tuberculosis (A15-A19)	1	982	5,0
Human immunodeficiency virus [HIV] disease (B20-B24)	2	666	3,4
Influenza and pneumonia (J09-J18)	3	653	3,3
Other viral diseases (B25-B34)	4	455	2,3
Episodic and paroxysmal disorders (G40-G47)	5	376	1,9
Other forms of heart disease (I30-I52)	6	256	1,3
Cerebral palsy and other paralytic syndromes (G80-G83)	7	245	1,2
Chronic lower respiratory diseases (J40-J47)	8	185	0,9
Inflammatory diseases of the central nervous system (G00-G09)	8	182	0,9
Cerebrovascular diseases (I60-I69)	8	179	0,9
Other Natural		6 072	30,9
Non-natural		9 427	47,9
All Causes		19 678	100,0

* Including deaths due to *MDR-TB* and *XDR-TB*

4.6.6 Leading underlying natural causes of death by province of death occurrence

Table 4.10 shows the ten leading underlying causes of death by province of death occurrence. Deaths that occurred outside South Africa and those where province of death occurrence was not specified in the death notification form, are not included in the table.

Diabetes mellitus was the leading cause of death in five provinces. In Western Cape (7,3%), KwaZulu-Natal (7,0%), Limpopo (6,9%), Eastern Cape (5,8%) and Gauteng (4,6%). It was the second leading underlying cause of death in Mpumalanga (6,3%), Free State (5,9%) and North West (5,7%). It was the third leading underlying cause in Northern Cape (5,4%).

Hypertensive diseases was the leading underlying cause of death in North West (8,5%), Northern Cape (8,0%), Free State (7,1%) and Mpumalanga (6,4%). Six underlying causes of death were common in all nine provinces, namely *diabetes mellitus*, *cerebrovascular diseases*, *hypertensive diseases*, *HIV disease*, *tuberculosis* and *other forms of heart diseases*.

However, the ranks of these causes of death differed between provinces. For example, *cerebrovascular diseases* ranked second in Western Cape and Kwa-Zulu Natal, third in Eastern Cape, Free State, North West, Gauteng and Limpopo.

Western Cape was the only province where *malignant neoplasms of respiratory and intrathoracic organs* was in the top ten leading underlying causes of death. It ranked ninth responsible for 3,5% deaths. This was the case with *intestinal infectious diseases*, it was amongst the top ten leading underlying causes of death only in Limpopo, ranked tenth and was responsible for 2,0% deaths in the province.

The underlying causes of death for all the provinces were a combination of communicable and non-communicable diseases. Western Cape had the highest number of non-communicable diseases with eight out of the ten being non-communicable diseases, followed by Gauteng which had seven non-communicable diseases in the top ten. Eastern Cape, Northern Cape, Free State, KwaZulu-Natal, North West and Mpumalanga all had six non-communicable diseases in their top ten leading underlying causes.

Table 4.10 - The ten leading underlying natural causes of death in each province of death occurrence, 2024

Causes of death (ICD-10)	Western Cape			Eastern Cape			Northern Cape			Free State			KwaZulu-Natal			North West			Gauteng			Mpumalanga			Limpopo		
	Rank	No.	%	Rank	No.	%	Rank	No.	%	Rank	No.	%	Rank	No.	%	Rank	No.	%	Rank	No.	%	Rank	No.	%	Rank	No.	%
Diabetes mellitus (E10-E14)	1	4 230	7,3	1	4 450	5,8	3	783	5,4	2	1 678	5,9	1	6 235	7,0	2	1 768	5,7	1	4 615	4,6	2	1 920	6,3	1	3 059	6,9
Cerebrovascular diseases (I60-I69)	2	3 800	6,6	3	4 284	5,5	4	735	5,0	3	1 532	5,4	2	5 577	6,3	3	1 409	4,6	3	4 288	4,3	4	1 510	5,0	3	2 375	5,4
Hypertensive diseases (I10-I15)	3	3 022	5,2	4	4 066	5,3	1	1 168	8,0	1	2 011	7,1	3	3 967	4,5	1	2 625	8,5	2	4 328	4,3	1	1 960	6,4	2	2 804	6,3
Ischaemic heart diseases (I20-I25)	3	3 019	5,2	...	...	...	9	391	2,7	10	656	2,3	8	2 554	2,9	10	512	1,7	6	2 765	2,8	3	1 557	5,1	...	...	...
Human immunodeficiency virus [HIV] disease (B20-B24)	5	2 945	5,1	5	4 003	5,2	2	798	5,5	4	1 527	5,4	6	2 823	3,2	6	1 046	3,4	...	...	...	5	1 208	4,0	5	1 560	3,5
Tuberculosis (A15-A19)	6	2 510	4,3	2	4 334	5,6	5	716	4,9	6	1 052	3,7	4	3 730	4,2	4	1 291	4,2	7	2 543	2,5	7	1 103	3,6	6	1 379	3,1
Malignant neoplasms of digestive organs (C15-C26)	7	2 421	4,2	10	1 625	2,1	...	...	...	...	...	...	...	...	...	...	...	...	7	2 503	2,5	...	...	...	...	...	...
Chronic lower respiratory diseases (J40-J47)	8	2 349	4,1	6	2 486	3,2	6	570	3,9	9	721	2,5	...	...	...	9	749	2,4	...	...	...	...	...	...	...	...	...
Malignant neoplasms of respiratory and intrathoracic organs (C30-C39)	9	1 996	3,5	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
Other forms of heart disease (I30-I52)	10	1 664	2,9	7	1 959	2,5	10	298	2,0	7	984	3,5	5	2 826	3,2	7	1 006	3,3	5	3 205	3,2	9	893	2,9	7	1 197	2,7
Other viral diseases (B25-B34)	...	...	...	7	1 897	2,5	7	502	3,4	8	802	2,8	7	2 654	3,0	8	937	3,0	7	2 507	2,5	8	922	3,0	7	1 188	2,7
Influenza and pneumonia (J09-J18)	...	...	...	9	1 799	2,3	8	403	2,8	5	1 418	5,0	9	2 467	2,8	5	1 104	3,6	4	3 456	3,4	6	1 158	3,8	4	1 935	4,4
Renal failure (N17-N19)	...	...	...	...	...	...	...	...	...	...	...	...	10	1 870	2,1	...	...	...	10	2 384	2,4	10	721	2,4	9	1 008	2,3
Intestinal infectious diseases (A00-A09)	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	10	906	2,0	
Other Natural		22 068	38,2		35 373	45,8		6 665	45,6		12 927	45,4		41 258	46,4		14 771	47,9		53 455	53,3		13 589	44,5		22 629	51,0
Non-natural		7 733	13,4		11 039	14,3		1 581	10,8		3 188	11,2		13 041	14,7		3 608	11,7		14 280	14,2		3 962	13,0		4 313	9,7
All causes		57 757	100,0		77 315	100,0		14 610	100,0		28 496	100,0		89 002	100,0		30 826	100,0		100 329	100,0		30 503	100,0		44 353	100,0

* Including deaths due to *MDR-TB* and *XDR-TB*
 ... Category not in top ten

4.7 Non-natural causes of death

The focus of this subsection is on non-natural causes of death. Information on non-natural causes of death is important in South Africa, considering the high levels of violence experienced in the country. This section profiles non-natural causes of death based on all external causes of morbidity and mortality (V01-Y98) derived from the causes of death specified on the death notification forms.

On the death notification form, where insufficient details are provided to code the non-natural cause of death accurately, Stats SA codes such deaths as *other external causes of accidental injury or event of undetermined intent* in line with the recommendations of WHO in classifying unknown non-natural causes of death (WHO, 2009b). This therefore contributes to the high percentage of unspecified causes of non-natural deaths. Results on non-natural causes of death should therefore be interpreted carefully, by taking into account the fact that over three-quarters of non-natural causes of death were not adequately classified. The unexpected lower number of deaths due to *transport accidents, assault, complications of medical and surgical care, intentional self-harm or sequelae of external causes of morbidity and mortality* may have been partly the result of causes classified as *other external causes of accidental injury or event of undetermined intent*.

Just above thirteen percent (13,2%) of all deaths registered in 2024 (refer to Table 4.4) were due to *external causes of morbidity and mortality*. Table 4.11 below, shows the percentage distribution of broad groups of non-natural causes and the associated number of deaths. It is observed that the majority of non-natural causes of death resulted from *other external causes of accidental injury* (74,9%). In terms of all deaths, *other external causes of accidental injury* accounted for 9,9%. This group includes *discharge from other and unspecified firearms* as well as *other accidental hanging and strangulation*.

Assault was the second most common non-natural cause of death and accounted for 11,9% of non-natural causes and 1,6% of all reported deaths. The third most common cause of non-natural deaths was *transport accidents* at 6,2% and constituting 0,8% of all deaths, followed by *event of undetermined intent* (3,6%) and *complications of medical and surgical care* (2,6%). Deaths due to *intentional self-harm* and *sequelae of external causes of morbidity and mortality* contributed less than 1,0% of non-natural deaths combined.

Table 4.11 - Distribution of non-natural causes of death by broad groups, 2024

Causes of death (ICD-10)	Number	Percentage (%) of non-natural causes	Percentage (%) of all causes (N = 476 751)
Other external causes of accidental injury (W00-X59)	47 098	74,9	9,9
Assault (X85-Y09)	7 486	11,9	1,6
Transport accidents (V01-V99)	3 923	6,2	0,8
Event of undetermined intent (Y10-Y34)	2 265	3,6	0,5
Complications of medical and surgical care (Y40-Y84)	1 625	2,6	0,3
Intentional self-harm (X60-X84)	432	0,7	0,1
Sequelae of external causes of morbidity and mortality (Y85-Y89)	42	0,1	0,0
All non-natural	63 367	100,0	13,2


A breakdown of the 47 098 deaths due to *other external causes of accidental injury* identified in Table 4.11 is provided in Table 4.12 to provide information that can be used to better understand deaths due to this cause, which comprised nearly two-thirds of all non-natural deaths.

Table 4.12 below, shows that almost half of these deaths (44,1%) were due to *accidental exposure to other and unspecified factors*. This includes *exposure to unspecified factor causing fracture and exposure to other unspecified factors*. Majority of deaths in this group were from *exposure to an unspecified factor*. This was followed by deaths due to *exposure to inanimate mechanical forces* which were the second leading cause, responsible for 26,3% deaths in this group. This group includes *discharge from other and unspecified firearms* as well as *contact with knife or sword*. The third most common cause was *other accidental threats to breathing* (11,7%), which includes *accidental hanging and strangulation*. The fourth most commonly reported deaths due to other external causes of accidental injury was *exposure to smoke, fire and flames* (6,1%), followed by *accidental poisoning by and exposure to noxious substance* (6,0%).

Table 4.12 - Distribution of deaths due to other external causes of accidental injury, 2024

Causes of death (ICD-10)	Number	Percentage
Accidental exposure to other and unspecified factors (X58-X59)	20 774	44,1
Exposure to inanimate mechanical forces (W20-W49)	12 377	26,3
Other accidental threats to breathing (W75-W84)	5 520	11,7
Exposure to smoke, fire and flames (X00 - X09)	2 880	6,1
Accidental poisoning by and exposure to noxious substance (X40-X49)	2 837	6,0
Accidental drowning and submersion(W65-W74)	1 572	3,3
Exposure to electric current, radiation and extreme ambient air temperature and pressure (W85-W99)	556	1,2
Exposure to forces of nature (X30 - X39)	255	0,5
Falls (W00-W19)	205	0,4
Exposure to animate mechanical forces (W50-W64)	52	0,1
Contact with venomous animals and plants(X20-X29)	40	0,1
Contact with heat and hot substances(X10-X19)	21	0,0
Overexertion, travel and privation(X50-X59)	9	0,0
Total	47 098	100,0

Figure 4.7 below, shows the distribution of *other external causes of accidental injury* from 2020 to 2024. Across the period, the same pattern is observed, with *accidental exposure to other and unspecified factors*, *exposure to inanimate mechanical forces*, *other accidental threats to breathing*, and *exposure to smoke, fire and flames* consistently leading. *Accidental exposure to other and unspecified factors* increased substantially from 15 521 deaths in 2020 to a peak of 24 270 in 2022, before declining to 20 774 in 2024. Despite this decline, deaths remained above 20 000, representing an overall increase of 33,8% between the start and end of the observed period. *Exposure to inanimate mechanical forces* also showed a notable increase, rising from 8 439 deaths in 2020 to 12 377 in 2024, an increase of 46,7%. *Accidental poisoning and exposure to noxious substances* was the only cause that increased annually, rising from 937 in 2020 to 2 344 in 2021.

Figure 4.7 - Distribution of other external causes of accidental injury, 2020–2024*

*(1) Data for 2020–2023 have been updated with late registrations/delayed death notification forms processed in 2025/2026

4.7.1 Non-natural causes of death by age and sex

This subsection focuses on the distribution of non-natural causes of death by sex and broad age groups (0,1–14, 15–29, 30–44, 45–64 and 65+). For international comparison, age group 15–44 has been divided into two age groups (15–29 and 30–44) as recommended by the WHO (1992).

Table 4.13 shows the distribution of non-natural causes of death by sex and broad age groups for deaths that occurred in 2024. The absolute numbers and percentages for both sexes may not be similar to the results presented in Table 4.11, due to the exclusion of deaths with missing sex and age.

The first section of Table 4.13 showing both sexes indicates that for both sexes, the age group mostly affected by non-natural causes of death was age group 15–29, where 47,9% of all deaths in this age group were due to non-natural causes. The age group least affected by non-natural causes for both sexes was 65 years and older, where just 2,8% of deaths in this age group were due to non-natural causes. *Assault* was more common among those aged 15–29, accounting for 16,7% of non-natural deaths in this age group. *Complications of medical and surgical care* were highest amongst the elderly (13,3%).

Differentials by sex show higher proportions of non-natural deaths for males at 19,6% compared to 5,9% of female non-natural deaths. Moreover, for each of the age groups, males had higher proportions of deaths due to non-natural causes compared to females, with the gap much wider at age group 15–29 where as much as 59,7% of male deaths resulted from non-natural cause compared to 23,8% of females in the same

age group. This is the only age group where the proportion of non-natural deaths is more than that of natural deaths for males.

Comparison between male and female deaths due to non-natural causes show that the proportion of deaths due to *assault* were high for males (13,5%) compared to females (5,9%). For both sexes, non-natural deaths due to *complications of medical and surgical care* were higher at infancy (those aged less than a year) as well as among the elderly (those aged 65 years and older). However, this cause of death was high amongst females compared to males, with the proportion of female deaths due to *complications of medical and surgical care* at 5,9% compared to 1,7% for males.

The proportion of non-natural deaths due to *transport accidents* were higher amongst females (7,8%) compared to males (5,9%). For both sexes, *intentional self-harm* and *sequelae of external causes of morbidity and mortality* each comprised less than 1% of deaths. For all age groups, *other external cause of accidental injury* was the highest non-natural cause of death.

Table 4.13 - Underlying non-natural causes of death by age group and sex, 2024

Causes of death (ICD-10)	Number							Percentage (%)						
	0	1-14	15-29	30-44	45-64	65+	All ages	0	1-14	15-29	30-44	45-64	65+	All ages
Both sexes*														
Transport accidents (V01-V99)	13	278	966	1459	940	267	3923	1,8	7,9	5,4	6,3	7,9	4,8	6,2
Other external causes of accidental injury (W00-X59)	657	2964	12971	17317	9028	4161	47098	90,2	84,0	72,1	74,7	75,9	75,1	74,9
Intentional self-harm (X60-X84)	0	16	156	163	77	20	432	0,0	0,5	0,9	0,7	0,6	0,4	0,7
Assault (X85-Y09)	8	63	3009	3275	943	188	7486	1,1	1,8	16,7	14,1	7,9	3,4	11,9
Event of undetermined intent (Y10-Y34)	16	155	781	762	403	148	2265	2,2	4,4	4,3	3,3	3,4	2,7	3,6
Complications of medical and surgical care (Y40-Y84)	34	49	102	202	501	737	1625	4,7	1,4	0,6	0,9	4,2	13,3	2,6
Sequelae of external causes of morbidity and mortality (Y85-Y89)	0	2	6	7	9	18	42	0,0	0,1	0,0	0,0	0,1	0,3	0,1
Sub total	728	3 527	17 991	23 185	11 901	5 539	62 871	100,0	100,0	100,0	100,0	100,0	100,0	100,0
Non-natural causes	728	3 527	17 991	23 185	11 901	5 539	62 871	3,6	30,3	47,9	29,6	9,1	2,8	13,3
Natural causes	19 290	8 095	19 575	55 127	119 194	190 214	411 495	96,4	69,7	52,1	70,4	90,9	97,2	86,7
All causes	20 018	11 622	37 566	78 312	131 095	195 753	474 366	100,0	100,0	100,0	100,0	100,0	100,0	100,0
Males*														
Transport accidents (V01-V99)	7	166	720	1 149	696	168	2 906	1,7	7,8	4,8	5,8	7,5	5,5	5,9
Other external causes of accidental injury (W00-X59)	360	1 801	10 836	14 803	7 169	2 271	37 240	89,6	84,9	72,4	75,0	77,0	74,4	75,1
Intentional self-harm (X60-X84)	0	7	111	128	50	15	311	0,0	0,3	0,7	0,6	0,5	0,5	0,6
Assault (X85-Y09)	5	36	2 762	2 956	810	124	6 693	1,2	1,7	18,4	15,0	8,7	4,1	13,5
Event of undetermined intent (Y10-Y34)	10	85	484	595	308	84	1 566	2,5	4,0	3,2	3,0	3,3	2,8	3,2
Complications of medical and surgical care (Y40-Y84)	20	25	59	110	268	380	862	5,0	1,2	0,4	0,6	2,9	12,4	1,7
Sequelae of external causes of morbidity and mortality (Y85-Y89)	0	1	5	7	9	11	33	0,0	0,0	0,0	0,0	0,1	0,4	0,1
Sub total	402	2 121	14 977	19 748	9 310	3 053	49 611	100,0	100,0	100,0	100,0	100,0	100,0	100,0
Non-natural causes	402	2 121	14 977	19 748	9 310	3 053	49 611	3,8	33,2	59,7	39,1	12,3	3,6	19,6
Natural causes	10 273	4 262	10 126	30 777	66 274	81 729	203 441	96,2	66,8	40,3	60,9	87,7	96,4	80,4
All cause	10 675	6 383	25 103	50 525	75 584	84 782	253 052	100,0	100,0	100,0	100,0	100,0	100,0	100,0
Females*														
Transport accidents (V01-V99)	6	111	244	309	240	99	1 009	1,9	7,9	8,3	9,4	9,5	4,0	7,8
Other external causes of accidental injury (W00-X59)	291	1 156	2 076	2 402	1 800	1 882	9 607	90,9	82,7	70,6	72,7	71,4	76,0	74,1
Intentional self-harm (X60-X84)	0	9	45	35	27	5	121	0,0	0,6	1,5	1,1	1,1	0,2	0,9
Assault (X85-Y09)	3	27	241	306	130	63	770	0,9	1,9	8,2	9,3	5,2	2,5	5,9
Event of undetermined intent (Y10-Y34)	6	69	291	160	91	64	681	1,9	4,9	9,9	4,8	3,6	2,6	5,3
Complications of medical and surgical care (Y40-Y84)	14	24	43	91	233	357	762	4,4	1,7	1,5	2,8	9,2	14,4	5,9
Sequelae of external causes of morbidity and mortality (Y85-Y89)	0	1	1	0	0	7	9	0,0	0,1	0,0	0,0	0,0	0,3	0,1
Sub total	320	1 397	2 941	3 303	2 521	2 477	12 959	100,0	100,0	100,0	100,0	100,0	100,0	100,0
Non-natural causes	320	1 397	2 941	3 303	2 521	2 477	12 959	3,5	26,8	23,8	12,0	4,6	2,2	5,9
Natural causes	8 920	3 820	9 407	24 247	52 835	108 458	207 687	96,5	73,2	76,2	88,0	95,4	97,8	94,1
All causes	9 240	5 217	12 348	27 550	55 356	110 935	220 646	100,0	100,0	100,0	100,0	100,0	100,0	100,0

Excluding cases with unspecified age and sex.

4.7.2 Non-natural causes of death by province of death occurrence

The distribution of the non-natural causes of death by province of death occurrence for the year 2024 are depicted in Table 4.14. KwaZulu-Natal (14,6%), Eastern Cape (14,3%) and Gauteng (14,2%) had the highest proportion of deaths due to non-natural causes. The lowest percentage of deaths due to non-natural causes were observed in Limpopo (9,7%). KwaZulu-Natal has consistently been the province with the highest proportion of non-natural deaths in the previous years.

The most common cause of non-natural deaths in all provinces was *other external causes of accidental injury* where more than half of non-natural deaths resulted from this broad group in each province. The proportion of deaths due to *other external causes of accidental injury* was highest in Gauteng (82,3%), Mpumalanga (78,9%) and KwaZulu-Natal (75,6%) followed closely by Western Cape at 72,4%.

In 2024, Western Cape had the highest proportion of deaths due to *assault*, accounting for 17,5% of all non-natural deaths within the province. Eastern Cape trailed closely accounting for 16,3% non-natural deaths in the province. Deaths due to *assault* were lowest in Limpopo (7,7%) and Gauteng (8,0%).

Deaths due to *transport accidents* were highest in Limpopo responsible for 20,1% of non-natural deaths in the province followed by Northern Cape at 12,4% and North West at 12,0%. Historically, Limpopo and Northern Cape have always had the highest proportions of *transport accidents* in South Africa. Gauteng had the least proportion of deaths due to *transport accidents* at 0,9%. *Complications of medical and surgical care, intentional self-harm* and *sequelae of external causes of morbidity and mortality* were the least common, each contributing less than 4,0% of non-natural deaths in each province.

Table 4.14 - Underlying non-natural causes of death by province, 2024

Causes of death (ICD-10)	Western Cape		Eastern Cape		Northern Cape		Free State		KwaZulu-Natal		North West		Gauteng		Mpumalanga		Limpopo	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Transport accidents (V01-V99)	297	3,8	652	5,9	196	12,4	220	6,9	865	6,6	432	12,0	135	0,9	244	6,2	866	20,1
Other external causes of accidental injury (W00-X59)	5 601	72,4	8 114	73,5	969	61,3	2 350	73,7	9 865	75,6	2 526	70,0	11 750	82,3	3 125	78,9	2 700	62,6
Intentional self-harm (X60-X84)	39	0,5	62	0,6	57	3,6	12	0,4	127	1,0	19	0,5	63	0,4	019	0,5	33	0,8
Assault (X85-Y09)	1 350	17,5	1 799	16,3	243	15,4	408	12,8	1 499	11,5	346	9,6	1 148	8,0	357	9,0	330	7,7
Event of undetermined intent (Y10-Y34)	180	2,3	240	2,2	64	4,0	132	4,1	377	2,9	181	5,0	721	5,0	155	3,9	213	4,9
Complications of medical and surgical care (Y40-Y84)	261	3,4	166	1,5	051	3,2	062	1,9	296	2,3	102	2,8	454	3,2	061	1,5	169	3,9
Sequelae of external causes of morbidity and mortality (Y85-Y89)	5	0,1	6	0,1	1,0	0,1	4	0,1	12	0,1	2	0,1	9	0,1	1,0	0,0	2	0,0
Total	7 733	100,0	11 039	100,0	1 581	100,0	3 188	100,0	13 041	100,0	3 608	100,0	14 280	100,0	3 962	100,0	4 313	100,0
Non-Natural causes	7 733	13,4	11 039	14,3	1 581	10,8	3 188	11,2	13 041	14,6	3 608	11,7	14 280	14,2	3 962	13,0	4 313	9,7
Natural causes	50 024	86,6	66 276	85,7	13 029	89,2	25 308	88,8	75 961	85,4	27 218	88,3	86 049	85,8	26 541	87,0	40 040	90,3
Total	57 757	100,0	77 315	100,0	14 610	100,0	28 496	100,0	89 002	100,0	30 826	100,0	100 329	100,0	30 503	100,0	44 353	100,0

*Excluding deaths that occurred outside South Africa and deaths with unspecified province of death.

4.8 Comparison between immediate, contributing and underlying causes of death

This subsection provides information on the total number of causes of death reported on each form. As previously mentioned in section 4.2, the death notification form provides for the recording of multiple causes of death. Section G of both death notification forms (BI-1663 and DHA-1663) makes provision for several causes to be reported on the form (see Appendix B). A maximum number of six causes can be recorded on the death notification form. These causes are recorded as immediate, contributing or underlying causes of death (see definitions in Appendix A).

This subsection provides information on the total number of causes of death entered on each form when reporting the cause of death. The total number of causes mentioned on each form were aggregated and grouped by broad groups of underlying causes of death and were then ranked.

The 20 leading causes based on all causes of death recorded on each form are shown in Table 4.15. The list includes both natural and non-natural causes, as well as deaths due to symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified to indicate the frequency of mentioning any cause on the death notification form.

Other forms of heart diseases were the most frequently recorded cause of death in 2024, mentioned in 73 672 (14,8%) death notification forms, followed by *hypertensive diseases* mentioned in 67 616 (13,6%) forms. *Influenza and pneumonia* was the third most recorded cause of death on 37 616 (7,6%) death notification forms.

Table 4.15 - Distribution of the 20 most reported causes of death, 2024

Rank	Causes of death (ICD-10)	Number of deaths in which the causes was reported	Percentage (%) of all deaths
7	Other forms of heart disease (I30-I52)	73 672	14,8
2	Hypertensive diseases (I10-I15)	67 616	13,6
6	Influenza and pneumonia (J09-J18)	37 616	7,6
1	Diabetes mellitus (E10-E14)	36 001	7,2
12	Renal failure (N17-N19)	35 488	7,1
3	Cerebrovascular diseases (I60-I69)	35 279	7,1
13	Other bacterial diseases (A30-A49)	32 572	6,5
4	Tuberculosis (A15-A19)	30 969	6,2
8	Ischaemic heart diseases (I20-I25)	20 155	4,0
9	Chronic lower respiratory diseases (J40-J47)	19 494	3,9
10	Other viral diseases (B25-B34)	19 069	3,8
5	Human immunodeficiency virus [HIV] disease (B20-B24)	18 926	3,8
17	Malignant neoplasms of ill-defined, secondary and unspecified sites (C76-C80)	14 583	2,9
19	Other acute lower respiratory infections (J20-J22)	11 666	2,3
11	Malignant neoplasms of digestive organs (C15-C26)	11 114	2,2
16	Intestinal infectious diseases (A00-A09)	10 796	2,2
14	Malignant neoplasms of female genital organs (C51-C58)	7 028	1,4
15	Malignant neoplasms of respiratory and intrathoracic organs (C30-C39)	6 319	1,3
18	Malignant neoplasms of breast (C50)	4 805	1,0
20	Malignant neoplasms of male genital organs (C60-C63)	4 625	0,9

*Including deaths due to *MDR-TB* and *XDR-TB*

The ten leading underlying natural causes of death shown in Table 4.5 are presented in Table 4.16 to show the breakdown of the number of deaths by whether the cause of death was reported as the immediate or contributing cause and selected as the underlying cause or whether it was selected only as the underlying cause but not mentioned as a cause on the form. The latter means the deceased died from a consequence of a disease without that specific disease being mentioned as an immediate or contributing cause.

Within each category, the counts of underlying causes and contributing causes are not duplicated, so that they can be summed up to equal the total number of times a specific cause of death was mentioned in the 2024 deaths. For example, 28 174 deaths had *diabetes mellitus* as the underlying cause and mentioned on the form as a contributing cause of death while 593 deaths had *diabetes mellitus* as an underlying cause but was not mentioned as a contributing cause on the form. This gives a total of 28 767 times *diabetes mellitus* was mentioned on the forms. In eight of the ten leading causes of death, 99% of the time where a contributing cause was mentioned, it was selected as the underlying cause of death as well. The exception was on *Hypertensive diseases* and *Human immunodeficiency virus [HIV] disease*. *Human immunodeficiency virus [HIV] disease* was mentioned as an underlying cause a total of 18 025 times, however, in 86,1% of those cases, it was not mentioned anywhere on the death notification form as a contributing cause of death.

Table 4.16 - Number and percentage (%) of deaths selected as underlying or reported as immediate or contributing causes of death, 2024

Causes of death (ICD-10)	Underlying rank	Number of deaths			Percentage (%) of any mention		
		Underlying and contributing cause	Underlying and not contributing cause	Total	Underlying and contributing cause	Underlying and not contributing cause	Total
Diabetes mellitus (E10-E14)	1	28 174	593	28 767	97,9	2,1	100,0
Hypertensive diseases (I10-I15)	2	16 756	9 216	25 972	64,5	35,5	100,0
Cerebrovascular diseases (I60-I69)	3	25 414	125	25 539	99,5	0,5	100,0
Tuberculosis (A15-A19)	4	18 598	68	18 666	99,6	0,4	100,0
Human immunodeficiency virus [HIV] disease (B20-B24)	5	2 505	15 520	18 025	13,9	86,1	100,0
Influenza and pneumonia (J09-J18)	6	14 945	14	14 959	99,9	0,1	100,0
Other forms of heart disease (I30-I52)	7	14 049	10	14 059	99,9	0,1	100,0
Ischaemic heart diseases (I20-I25)	8	13 256	95	13 351	99,3	0,7	100,0
Chronic lower respiratory diseases (J40-J47)	9	12 070	11	12 081	99,9	0,1	100,0
Other viral diseases (B25-B34)	10	11 929	16	11 945	99,9	0,1	100,0

*Including deaths due to *MDR-TB* and *XDR-TB*.

5. Conclusion

Registering causes of death is essential for producing reliable mortality statistics, which inform public health policies, enable disease surveillance and guide resource allocation. Accurate data helps authorities identify leading causes of mortality (e.g. communicable diseases, noncommunicable, or injuries) and track trends over time. This allows for effective health planning, such as allocating resources for specific diseases or targeting interventions to reduce premature deaths. Where this information is well captured, it provides timely information for implementation of interventions during health emergencies as well as preventive and curative measures within the health system. The underlying causes of deaths provide data on leading causes of deaths in the population based on civil registration.

This statistical release provides information on mortality and causes of death for deaths that occurred and were registered in South Africa in 2024 as well as registered deaths of previous years that reached Stats SA during the 2025/2026 processing phase. After a protracted decline in mortality over several years, increases of 17,1% between 2020 and 2021 were observed due to the emergence of COVID-19. However, deaths that occurred from 2022 show a reversal of the pattern seen in 2020 and 2021. The number of deaths decreased from 624 528 in 2021 to 493 900 in 2022 and further declined to 474 366 in 2024.

In 2024, the proportion of male to female deaths was almost the same from infancy until age group 15–19 after which there were more male deaths than female deaths. This trend reversed from age group 70 years and older where there were more female deaths than male deaths. The highest number of deaths amongst females were aged 70–74 (9,5%) and for males it was those aged 60–64 at 9,2%. The lowest number was observed among those aged 5–9 for both sexes 5–9 at 0,6% for males and 0,5% for females. The distribution of deaths by population group show that black Africans had the highest percentage of deaths followed by the white, coloured and Indian/Asian population groups. However, computation of race specific death rates show that white population group had the highest death rate followed by the coloured, Indian/Asian and black African population groups.

Geographic distribution of deaths by province of death occurrence shows that the highest proportion of deaths (21,2%) occurred in Gauteng province, followed by KwaZulu-Natal (18,8%) and then Eastern Cape (16,3%). The lowest proportion of deaths occurred in Northern Cape (3,1%). The highest proportion of deaths occurred in hospitals (40,6%) followed by deaths occurring at home (23,3%).

From 2003 to 2024, communicable diseases gradually decreased from 45,6% to 26,0%, with a noticeable increase during the COVID-19 pandemic. In contrast, non-communicable diseases increased from 44,9% to 60,7% over the same period. Non-natural deaths also increased, from accounting for 9,5% of all deaths in 2003 to 13,3% in 2024.

The top three causes of death in 2024 were due to non-communicable diseases, with *diabetes mellitus* accounting for 6,1% deaths, followed by *hypertensive diseases* and *cerebrovascular diseases* at 5,5% and 5,4% respectively. *Tuberculosis* was the leading cause of death amongst males accounting for 4,8% deaths while *diabetes mellitus* was the leading cause of death amongst females responsible for 7,9% deaths. For both sexes, six of the 10 leading causes of death were non-communicable diseases. The cause of death on the top

10 leading causes of death for males but not for females was *chronic lower respiratory diseases*. While the opposite for women was *malignant neoplasms of female genital organs*.

Looking at leading causes of deaths by broad age groups, *influenza and pneumonia* were in the top ten leading causes of death for all age groups. *Tuberculosis* and *other forms of heart diseases* were among the 10 leading underlying causes of death in most age groups, excluding infants and persons aged 65 years and older, while *cerebrovascular diseases* and *mellitus diseases* were among the ten leading underlying causes of death for from age 15 years and older. The leading underlying cause of death for infants (age 0) was *respiratory and cardiovascular disorders specific to the perinatal period*, responsible for 14,9% of deaths. *Influenza and pneumonia* were the leading underlying cause of death for age group 1–14 years, accounting for 5,5% of deaths. *Human immunodeficiency virus (HIV)*, and *tuberculosis* were the two leading underlying cause of death among those aged 15–44 accounting for 7,7% and 7,0% of deaths respectively. For age group 45–64 years, *diabetes mellitus* was the leading cause of death responsible for 7,1% deaths and finally *hypertensive diseases* were the leading cause of death for those aged 65 years and older at 9,4%.

Diabetes mellitus, *chronic lower respiratory diseases*, *hypertensive diseases*, *tuberculosis* and *other forms of heart diseases* were in the top 10 leading causes of death for all the provinces. *Diabetes mellitus* was the leading cause of death in five of the nine provinces. It was not the leading cause of death in Northern Cape, Free State, North West and Mpumalanga. The leading cause of death in Northern Cape, Free State, North West and Mpumalanga was *hypertensive diseases*.

As expected, males had a higher proportion of deaths attributed to non-natural causes at 19,6% compared to females at 5,9%. Males aged 15–29 years remained the group most burdened by deaths due to non-natural causes (59,7%). *Transport accidents* had a higher proportion of female deaths at 7,8% compared to males at 5,9%. *Complications of medical and surgical care* were highest amongst females particularly at age 65 years and older (14,4%)

Although improvements have been made in the reporting of causes of death and completeness of death registration, further improvements are needed in the reduction of unspecified and unknown cases. This applies for both the natural as well as non-natural causes. For natural causes, *symptoms and signs not elsewhere classified* accounted for 18,2% deaths that occurred in 2024. High levels of ill-defined causes make it impossible to monitor, evaluate, or compare disease trends over time. It also creates a distorted view of the health status of a population, often masking the true, leading causes of death.

References

AbouZahr C, de Savigny D (2017). Towards universal civil registration and vital statistics systems: the time is now. *Lancet*. 2017;390(10091):1407-1408.

Bennett, N.G. and Horiuchi, S. (1981). Estimating the Completeness of Death Registration in a Closed Population. *Population Index*, 47(2): 207–21.

Bennett, N.G. and Horiuchi, S. (1984). Mortality estimation from registered deaths in less developed countries. *Demography*, 21(2): 217–234.

Hill, K. (1987). Estimating Census and Death Registration Completeness. *Asian and Pacific Population Forum*, 1(3): 8–13, 23–24.

Institute for Health Metrics and Evaluation. <https://www.healthdata.org/>. Accessed 01 August 2025

Mahapatra, P., Shibuya, K., Lopez, A., Coullare, F., Notzon, F.C. and Szreter, S. On behalf of the Monitoring Vital Events (MoVE) writing group. (2007). Civil registration systems and vital statistics: successes and missed opportunities. *The Lancet*, 370 (10): 1653–1663.

National Planning Commission (2012) National Development Plan 2030: Our future – make it work. Pretoria: The Presidency, Republic of South Africa. Available at: <https://www.gov.za/documents/national-development-plan-2030-our-future-make-it-work>

Peters G.B. (2016). Civil Registration and Vital Statistics as a Tool to Improve Public Management. Department of Political Science, University of Pittsburgh. Institutions for Development Sector. Downloaded from <https://publications.iadb.org/english/document> on 04/08/2025.

Republic of South Africa. (2014). Regulations on the Registration of Births and Deaths (Government notice No. 37373). Pretoria: Government of South Africa.

Stats SA (Statistics South Africa). (2020). Mid-year population estimates, 2020 (P0302). Pretoria: Statistics South Africa.

Stats SA (Statistics South Africa). (2021). Mortality and causes of death in South Africa, 2018: findings from death notification form (P0309.3). Pretoria: Statistics South Africa

Stats SA (Statistics South Africa). (2023). Mortality and causes of death in South Africa, 2019: findings from death notification form (P0309.3). Pretoria: Statistics South Africa.

Statistics South Africa (2024) Statistics Amendment Act, 2024 (Act No. 29 of 2024). Pretoria: Government Printer. Available at: https://www.statssa.gov.za/?page_id=18225

UN (United Nations). (2014). Principles and recommendations for a vital statistics system, Third Revision. Department of economic and social affairs: United Nations publication.

WHO (World Health Organization). (2013). Strengthening civil registration and vital statistics for births, deaths and causes of death: resource kit. Geneva: World Health Organization.

WHO (World Health Organization). (2014a). Analysing mortality levels and causes-of-death (ANACoD) Electronic Tool, Version 2.0. Department of Health Statistics and Information Systems. Geneva: World Health Organization.

WHO (World Health Organization). (2014b). Performing routine basic checks on compiling cause-of-death data (CoDEdit) Electronic Tool, Version 1.0. Department of Health Statistics and Information Systems. Geneva: World Health Organization.

WHO (World Health Organization). (2016). International classification of diseases and related health problems. Tenth Revision. Volume 2: 2016 Edition. Geneva: World Health Organization.

WHO (World Health Organization). (2019) ICD-11 Implementation or Transition Guide. Geneva.

World health statistics overview 2019: monitoring health for the SDGs, sustainable development goals. Geneva: World Health Organization; 2019.

"World Bank; World Health Organization. 2014. Global Civil Registration and Vital Statistics: A Scaling Up Investment Plan 2015-2024. © <http://hdl.handle.net/10986/18962> License: CC BY 3.0 IGO."

Appendices

Appendix A: Glossary

Causes of death are all those diseases, morbid conditions, or injuries that either resulted in or contributed to death, and the circumstances of the accident or violence which produced any such injuries.

Contributing causes of death are morbid conditions, if any, giving rise to the immediate cause of death.

COVID-19 a highly contagious infectious disease caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2)

Death is a permanent disappearance of all evidence of life at any time after a *live birth* has taken place.

Human immunodeficiency virus (HIV) is the pathogenic organism responsible for the acquired immunodeficiency syndrome (AIDS), also known as the lymphadenopathy virus (LAV).

Immediate cause of death is the disease or condition directly leading to death.

Leading underlying causes of death are the most frequent underlying causes of death in any given population. In this release, the underlying causes of death are ranked according to frequency.

Live birth in relation to a child, means the birth of a child born alive.

Multiple causes of death are all morbid conditions, diseases and injuries entered on the death certificate. These include those involved in the morbid train of events leading to the death which were classified as either the underlying cause, the intermediate cause, or any intervening cause and those conditions which contributed to death but were not related to the disease or condition causing death.

Neonatal death is the death of a live-born child during the first 28 completed days of life.

Perinatal deaths are a combination of stillbirths and infants who die in the first week after birth (early neonatal deaths).

Post-neonatal death is a live-born infant dying after 28 completed days of birth but before the first year of life is completed.

Population group: according to the Population Registration Act Repeal Act (Act No. 114 of 1991), the South African Population Register no longer stores information regarding the population group of individuals whose details are on the register. This Repeal Act is still in place; therefore, the population group used in this report

refers to the population group as identified by the certifying physician/professional nurse on the death notification form and is only used for statistical purposes.

Stillbirth is the intra-uterine death of a foetus of at least 26 weeks of gestation that showed no sign of life after complete birth.

Underlying cause of death (previously known as primary cause) is the disease or injury that initiated the sequence of events leading directly to death; or the circumstances of the accident or violence which produced the fatal injury.

Appendix B: Death Notification form

Please refer to the Mortality and causes of death in South Africa: Findings from death notification, 2016 on pages 59–64 for copies of both the BI–1663 and DHA–1663 forms (Stats SA, 2018).

Appendix C: Assessment of the quality of data

The gold standard in mortality statistics is to have real-time data on the number of deaths and corresponding medically certified causes of death (WHO, 2013). However, the information needs to be of the highest quality in terms of completeness of death registration, timeliness of death registration and publication of death statistics, and accuracy of information provided embedded in deaths with correct information on characteristics of deceased, accurate causes of deaths and lower proportions of deaths with ill-defined or unspecified causes of deaths. In this regard, data quality confrontation has to be undertaken for improvements in mortality statistics to be realised. Improvements in quality of mortality data are essential in more effective policies and programmes concerning people's health and quality of life with the aim of leaving no one behind.

Completeness of death registration

The proportion of all deaths that occurred in a specific period and were covered by the civil registration of a country (referred to as completeness) was estimated at 96% for adult deaths (15 years and older) for the intercensal period 2011–2016. The completeness level for male adult deaths was estimated at 97% whereas for females it was slightly lower at 95%. The methods used to derive the level of completeness for the intercensal period 2011–2016 were the Generalised Growth Balance (GGB) as proposed by Hill (1987), and the Synthetic Extinct Generation method (SEG) by Bennett and Horiuchi (1981, 1984). For the underlying assumptions and method followed, refer to Stats SA (2014). The extent of completeness of child deaths registration (0–14 years) is less certain, given the lack of completeness level estimates. For the 2024 adult death registration, the 96% completeness level is adopted

Timeliness of death registration

In South Africa, the Regulations for the Registration of Births and Deaths published in 2014 mandate that a death must be registered within 72 hours (3 days) of occurrence (Republic of South Africa, 2014). Timeliness in death registration indicates that all deaths are registered within the legally stipulated time allowance (UN, 2014). In general, timeliness of death registration refers to the interval between the date of death occurrence and the date it was registered with the Department of Home Affairs (DHA).

The number of days it took for deaths to be registered at DHA offices in 2024 is shown in Table C.1. For deaths that occurred in 2024, 15,9% were registered within a day of occurrence, 30,6% a day after the death had occurred, 18,3% on the second day after death occurrence and 12,9% on the third day. The proportion of deaths which were registered within the 72 hours (3 days) stipulated by the regulations was 77,7%.

Table C.1 – Distribution of deaths by the number of days it took to register the death, 2024

Number of days	Number of deaths	Percentage (%)	Cumulative percentage (%)
Within a day of death	75 565	15,9	15,9
1 day	144 980	30,6	46,5
2 days	86 705	18,3	64,8
3 days	61 206	12,9	77,7
4 days	35 524	7,5	85,2
5 days	20 572	4,3	89,5
6 days	12 152	2,6	92,1
7-13 days	21 906	4,6	96,7
14-20 days	3 607	0,8	97,4
21-30 days	2 179	0,5	97,9
31-364 days	9 617	2,0	99,9
1 year+	353	0,1	100,0
Total	474 366	100,0	

Timeliness of publication of statistics

The United Nations (UN) recommends that a one-year time lapse from the end of the reference period to publication and dissemination of death statistics from the civil registration be maintained in order for vital statistics to be considered timely (UN, 2014). This statistical release fell short of this recommendation as it is published 21 months from the time the event occurred. Since civil registration deaths are continuously updated, the proportion of total registrations that are delayed or late provide an estimate of under-reporting in previous time periods.

Table C.2 shows the number of deaths published in the 2023 mortality and causes of death report for the years 2003 to 2023, and late or delayed death registrations processed during the 2025/2026 processing year. The table shows that 14 851 additional death notification forms for deaths that occurred between 2003 and 2023 were processed during the 2025/2026 processing phase. Majority (6 693) of the additional forms were for deaths that occurred in 2023. The distribution of deaths for 2003 to 2023 updated for late or delayed death notification forms is provided in Appendix D.

Table C.2 – Number of deaths published in 2023 publication and late registrations processed during the 2025/2026 processing phase by year of death, 2003–2023

Year of death	Number of deaths published in 2023 publication	Additional forms received in the 2025/2026 processing phase	Total number of deaths
2003	559 161	78	559 239
2004	579 069	111	579 180
2005	600 236	70	600 306
2006	614 782	82	614 864
2007	606 813	69	606 882
2008	598 856	76	598 932
2009	584 652	66	584 718
2010	553 395	85	553 480
2011	520 812	195	521 007
2012	498 278	266	498 544
2013	479 805	810	480 615
2014	480 166	272	480 438
2015	478 220	454	478 674
2016	475 288	264	475 552
2017	470 123	470	470 593
2018	471 649	598	472 247
2019	468 183	640	468 823
2020	517 120	655	517 775
2021	623 132	1 396	624 528
2022	492 399	1 501	493 900
2023	476 605	6 693	483 298
Total	11 148 744	14 851	11 163 595

Data confrontation

In previous years, as part of data quality assessment, data confrontation was undertaken with other data sources, primarily, mortality data in the National Population Register data (NPR). However, in the mortality and causes of death 2024 statistical release this process was not undertaken as was case in previous publications. This was attributable to Stats SA not having access to mortality data captured in the national population register at DHA. As a result, the section dealing with comparison of the two datasets in order to assess coherency is excluded. It is envisaged that in forthcoming publications the assessment will be performed.


Quality of causes of death information

Quality information on the underlying causes of death is critical to guide decision-making in public health. As such, it is important that this information is assessed from data processing through to the data analysis phase in order to measure the extent to which the data may be used for health policies and programmes. The ill-defined causes, typically classified under ICD-10 codes R00–R99, refer to diagnoses that are vague, non-specific and have insufficient details to be of value in promoting preventive and curative health interventions. Although ill-defined causes still help to provide the overall mortality due to broad diseases, they fail to provide a concise picture as they poorly attribute the underlying cause and therefore, widely regarded as an indicator of poor data quality.

Ill-defined causes of death, typically classified under ICD-10 codes R00–R99, remain a persistent challenge in mortality and cause-of-death data in South Africa. This group is widely regarded as an indicator of poor data quality. In South Africa, these causes ranged from 12,7% to 13,6% from 2003 to 2018. From 2019–2024, an

increase in proportion of deaths due to this group, from 15,7% to 18,2%, was observed. The 2019 increase coincides with the implementation of a WHO recommendation to code deaths due to cardiac arrest (I46) as ill-specified (R99), marginally inflating the number of ill-specified causes.

Figure C 2 – Percentage distribution of deaths assigned to symptoms, signs and abnormal clinical and laboratory findings not elsewhere classified and year of death 2003-2024*



*Data for 2003–2024 have been updated with late registrations/delayed death notification forms processed in 2025/2026.

Assessment framework for death registration data

This statistical release adopts the assessment framework proposed by Mahapatra et al. (2007) to assess the quality of the 2024 death registration data received from the civil registration system. The framework proposed two categories, namely general vital statistics and causes-of-death statistics. Both categories measure quality in terms of level of accuracy, relevance, comparability, timeliness and accessibility.

The results of the Mahapatra et al. 2007 assessment framework for the 2024 mortality and causes of death data from the South African civil registration system are shown in Table C.3 and Table C.4. Table C.3 shows the percentage of key variables with unknown or unspecified information and forms part of the accuracy criteria in the assessment framework for the year 2024. The unknown cases refer to cases where more than one option was selected on the form or where the information could not be classified according to specified categories while unspecified cases refer to missing data for that variable.

Table C.3 shows that 0,1% of deaths had unknown/unspecified sex, while 0,0% of deaths had unknown/unspecified age of deceased and province of death occurrence. Unknown/unspecified province of

usual residence was 0,7% in 2024. The information on province of birth occurrence remained relatively the same at 23,7% in 2023 to 23,6% in 2024. Missing information for marital status was 27,5%.

The results further indicate that 15,0% of the deaths had unknown or unspecified information on population group. This variable has been improving in the recent years considering that over the period 1997 to 2014 missing information on this variable was constant at around 25%.

In this release, no analyses were undertaken for all variables where almost half or more of the deaths had unknown or unspecified information. In 2024, education level (52,3%); industry (88,2%) and pregnancy status (79,4%) remained the variables with almost half or more of the information classified as unknown or unspecified. However, a dataset containing unit records on mortality and causes of death 2024, which include variables not covered in this release due to poor reporting, is available on request from Stats SA.

Table C.3 – Percentage (%) of deaths classified as unknown/unspecified for selected variables, 2024

Variables	Applicable group	Percentage (%) unknown or unspecified
Sex	All	0,1
Age	All	0,0
Province of death occurrence	All	0,0
Province of usual residence of deceased	All	0,7
Province of birth	All	23,6
Population group	All	15,0
Place or institution of death occurrence	All	28,7
Method used to ascertain cause of death	All	2,3
Marital Status	All	27,5
Smoking status	Aged 16 and older	32,2
Education	Aged 5 and older	52,3
Occupation	Aged 15 and older	2,8
Industry	Aged 15 and older (economically active)	88,2
Pregnancy status	Females aged 10–55	79,4

In addition, for the accuracy dimension, Table C.4 indicates that 96% of adult (15 years and older) death registrations were reported for the 2011–2016 intercensal period. The table also shows that the relevance and comparability of general vital statistics is regarded as complete.

The table further shows that for causes-of-death-statistics, 44% of the 2024 deaths occurred within a health-care facility. This approximates the percentage of deaths whose causes are more likely to be detailed enough for the underlying cause to be derived. Cause-of-death-statistics are regarded as completely relevant as they are based on routine tabulations by sex and five-year age groups as well as the fact that tabulation of cause-of-death information is provided for the nine provinces and 52 district municipalities in the country. The tools used in coding causes of death (International Classification of Diseases 10th revision) for 2024, and the variables analysed were similar to those in previous years. Therefore, comparability over time and with other countries is also regarded as complete.

Table C.4 – Assessment of the 2024 South African death statistics from civil registration system using the framework proposed by Mahapatra et al. (2007)

General vital statistics		Cause-of-death statistics	
Criteria and indicators	Measure	Criteria and indicators	Measure
Accuracy Completeness of death registration	96%	Accuracy Proportion of deaths that occurred in healthcare facilities	44,0%
Missing data	See Appendix C - Table C3	Proportion of deaths assigned to symptoms and signs of disease not elsewhere classified	18,2%
Relevance Routine tabulations by sex and 5-year age groups	100%	Relevance Routine tabulation by sex and 5-year age groups	100%
Deaths in children under five years tabulated by 0 and 1-4-year age group	100%	Number of cause-of-death tabulation areas	9 provinces and 52 district municipalities
Comparability Stability of key definitions over time	100%	Comparability Consistency of cause specific mortality proportions over consecutive years	100%
Uniformity of definitions across areas	100%	ICD coding for certification and coding of causes of death, revision used and code level to which tabulations are published	Coding causes of death using the 10th revision at 3/4-digit level
Timeliness Processing time	10 months		
Mean time from end of reference period to publication	21 months		
Accessibility Media - number of formats in which data are released	Data portal on website		
Metadata	Data portal on the website and available on request		
Availability of user service	Email: info@statssa.gov.za / Tel: 012 310 8600 / Fax (012) 310 8500 / 8495		